


UNIVERSIDADE FEDERAL DO PARANÁ

ANNA CLAUDIA BAUMEL MONGRUEL



MYCOPLASMA SP. IN SHEEP FROM SOUTHERN BRAZIL

CURITIBA

2019

ANNA CLAUDIA BAUMEL MONGRUEL

Mycoplasma sp. in sheep from southern Brazil

Dissertação apresentada ao curso de Pós-Graduação em Ciências Veterinárias, Setor de Ciências Agrárias, Universidade Federal do Paraná, como requisito à obtenção do título de Mestre em Ciências Veterinárias.

Orientador: Prof. Dr. Rafael Felipe da Costa Vieira

Curitiba


2019

M743m Mongruel, Anna Claudia Baumel
Mycoplasma SP. in sheep from southern Brazil / Anna Claudia Baumel Mongruel. - Curitiba, 2019.
79 p.: il.,

Dissertação (Mestrado) - Universidade Federal do Paraná.
Setor de Ciências Agrárias, Programa de Pós-Graduação em Ciências Veterinárias.
Orientador: Rafael Felipe Da Costa Vieira

1. Ovino - Brasil. 2. Ovelha. 3. Anemia. I. Vieira, Rafael Felipe Da Costa (Orientador). II. Título. III. Universidade federal do Paraná.

CDU 636.3:616.155.194(816)



MINISTÉRIO DA EDUCAÇÃO
SETOR SETOR DE CIÊNCIAS AGRÁRIAS
UNIVERSIDADE FEDERAL DO PARANÁ
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO
PROGRAMA DE PÓS-GRADUAÇÃO CIÊNCIAS
VETERINÁRIAS - 40001016023P3


TERMO DE APROVAÇÃO


Os membros da Banca Examinadora designada pelo Colegiado do Programa de Pós-Graduação em CIÊNCIAS VETERINÁRIAS da Universidade Federal do Paraná foram convocados para realizar a arguição da dissertação de Mestrado de **ANNA CLAUDIA BAUMEL MONGRUEL** intitulada: *Mycoplasma sp. in sheep from southern Brazil*, após terem inquirido a aluna e realizado a avaliação do trabalho, são de parecer pela sua APROVAÇÃO no rito de defesa.

A outorga do título de mestre está sujeita à homologação pelo colegiado, ao atendimento de todas as indicações e correções solicitadas pela banca e ao pleno atendimento das demandas regimentais do Programa de Pós-Graduação.

CURITIBA, 20 de Maio de 2019.


RAFAEL FELIPE DA COSTA VIEIRA
Presidente da Banca Examinadora (UFPR)


IVAN ROQUE DE BARROS FILHO
Avaliador Interno (UFPR)


KATHERINNE MARIA SPERCOSKI
Avaliador Externo (UFPR/PALOT.)

AGRADECIMENTOS

Defender uma tese de mestrado sempre esteve nos meus planos profissionais. Mas, é lógico, planejar é sempre muito mais fácil do que executar. A execução desse plano não foi fácil, ou tranquila, da maneira que eu planejei. Mas ela aconteceu, muito por conta dos meus esforços e da minha vontade de não desistir, mas também pela participação indispensável de muitas outras pessoas. Vou começar agradecendo aos meus orientadores, Prof. Dr. Rafael Vieira e Dra. Thállitha Vieira. Graças a vocês, o crescimento profissional, e também pessoal, que foi obtido desde o meu primeiro dia como residente na Universidade Federal do Paraná até aqui, como mestranda dessa instituição, é imensurável. Agradeço os esforços, a paciência e as orientações, sempre prontamente disponibilizados.

Além dos meus orientadores, a equipe do laboratório com quem tive a oportunidade de trabalhar também foi indispensável para que eu chegasse até aqui. Obrigada especialmente à Jessica, Kelly, Carolina, Nayara, Viviane, Cássia e Maria. Cada uma de vocês colaborou para que esse trabalho acontecesse e também para que eu aprendesse muito nesses últimos anos. Obrigada ao pessoal da Universidade Estadual do Norte do Paraná, em especial aos professores Flávio Otomura e Liza Ogawa, pela parceria que permitiu a execução desse projeto.

Agradeço também aos meus pais, Roberto e Isabela, por me apoiarem de maneira irrestrita, deixando sempre claro o orgulho e satisfação que sentem pelas minhas escolhas e pela pessoa que eu sou (mesmo que isso envolva uma ou duas tatuagens). Agradeço aos meus irmãos: Felipe, Daniel, Cynthia e Marcello, por existirem e tornarem a existência mais fácil. Em especial, ao meu irmão Felipe, que foi uma das minhas principais companhias em Curitiba, sempre me deixando admirada e inspirada com a sua forma de pensar e encarar seus ideais. Em relação à minha família, preciso citar e agradecer o carinho dos meus avós Odeni e Zaclis. Se eu insisto tanto na carreira acadêmica, é para tentar ser pelo menos um pouco do que meu avô, como educador, é.

Se essa parte da minha vida em Curitiba trouxe algo que mudou minha vida pra sempre, foi ter conhecido o Guilherme (ou melhor, reconhecido). Guilherme, você presenciou de perto todos os meus surtos e tempestades em copo d'água e, mesmo assim, sempre foi paciente e motivador. Obrigada por todo o apoio e amor. E falando em pessoas que Curitiba me trouxe, não posso deixar de citar todos os amigos maravilhosos que conheci aqui: Daniel, Felipe, Luciana e Jonatas. Que alegria saber que o destino foi tão querido comigo, me trazendo pessoas com quem eu me dei tão bem. E devo citar as amigas de sempre, que também foram muito valiosas nessa caminhada: Salua, Nathália e Katherinne. Mesmo com a distância, e a falta de tempo, vocês sempre se demonstraram prontas para me ouvir e dizer as melhores palavras que alguém poderia falar, mesmo que fosse por mensagem.

Obrigada também à Antônia e ao Simbah, nossos bichinhos de estimação. Sempre que estou com eles, eu lembro do porquê eu escolhi cursar medicina veterinária. Enfim, obrigada a todos que, de alguma forma ou outra, me auxiliaram e torceram por esse dia.

“I don’t know where I’m going from here. But I promise it won’t be boring”

David Bowie

RESUMO

O rebanho mundial de ovinos é estimado em mais de um bilhão de animais e a América do Sul representa 6,8% da produção mundial de ovinos. Embora as ovelhas sejam criadas principalmente para fins de obtenção de carne, esses animais fornecem uma ampla gama de produtos, como leite, pele e lã. Em 2016, o rebanho ovino brasileiro atingiu o número estimado de 18.000.000 de animais somado a um crescente cenário de produção e consumo de produtos ovinos à longo prazo. O Paraná, localizado na região sul do país, possui um rebanho estimado em mais de 500.000 cabeças, representando 3,12% do rebanho nacional e considerado um negócio em crescimento. Diversos hemoparasitos, como bactérias e protozoários, são descritos infectando ovinos de todo o mundo, o que pode causar febre, anorexia, anemia e perdas econômicas envolvendo esses animais. O presente estudo tem como objetivo identificar o agente causador de um surto de anemia em um rebanho ovino do município de Bandeirantes, Paraná, sul do Brasil. Quarenta e seis ovinos mantidos no município de Bandeirantes, região norte do Paraná, foram avaliados molecularmente por PCR quanto à presença de hemoplasmas, piroplasmas e *Anaplasma* spp. A infecção por parasitas gastrintestinais foi avaliada pela contagem de ovos por grama (OPG). Coleta de carrapatos, avaliação hematológica e análise filogenética também foram realizadas. Todas as amostras foram negativas para a presença de material genético de *Anaplasma* spp. e piroplasmas. Sete amostras (16,6%) apresentaram ampliações do tamanho esperado dos *primers* utilizados para detecção de *Mycoplasma* spp. Três amostras foram escolhidas para sequenciamento e apresentaram similaridade de 99% com sequências dos fragmentos 16S rDNA de *M. ovis* depositados no banco de dados do GenBank. A análise de haplótipos demonstrou a presença de três genótipos diferentes. Dentre 38 animais avaliados, 24 (63,15%) apresentaram valores de EPG acima de 500 e foram considerados positivos para a presença de ovos do tipo Strongylida. Carrapatos da espécie *Rhipicephalus sanguineus* foram coletados de dois animais. Na análise filogenética, as sequências do presente estudo foram alocadas em um clado conjunto com sequências de *Mycoplasma* sp. e *Mycoplasma ovis* isoladas em hospedeiros de outras regiões do mundo. O presente estudo relata a ocorrência de *Mycoplasma* sp. em ovinos do estado do Paraná. A co-infecção de parasitas gastrointestinais e hemoplasmas pode agravar a condição de saúde de ovelhas, representando um alerta a cerca do diagnóstico de hemoplasmas em rebanhos de ovelhas onde ocorre infecções por parasitos gastrointestinais.

ABSTRACT

Worldwide sheep herd is estimated in more than one billion animals and South America represents 6.8% from the world sheep production. Although sheep are raised primarily for meat purposes, these animals provide a wide range of products, such as milk, skin and wool. In 2016, the Brazilian sheep herd reached the estimated number of almost 18,000,000 of animals added to an increasing long-term scenario of production and consumption of sheep products. Paraná State, located in the southern region, have an estimated herd of more than 500,000 sheep, representing 3,12% from the national herd and considered a growing business. A wide range of hemoparasites, such as bacteria and protozoa, are described infecting sheep from worldwide, which may cause febrile phases, anorexia, anemia and economic losses in sheep. The present study aims to identify the causative agent of an outbreak of anemia in a sheep herd from Bandeirantes County, Paraná State, southern Brazil. Forty-six sheep maintained at Bandeirantes county, north region of Paraná State, were molecularly evaluated by PCR for the presence of hemoplasmas, piroplasms and *Anaplasma* spp. Infection by gastrointestinal parasites were accessed by worm egg per gram counting (EPG). Collection of ticks, hematological evaluation and phylogenetic analysis were also performed. All samples were negative for the presence of genetic material from *Anaplasma* spp. and piroplasms. Seven samples (16,6%) obtained amplifications of the expected size from the primers used for *Mycoplasma* spp. detection. Three samples were choosing for sequencing and obtained a similarity of 99% with sequences of the 16S rDNA fragments from *M. ovis* deposited in GenBank database. Haplotype analysis showed the occurrence of three different genotypes. From 38 animals evaluated, 24 (63,15%) presented EPG values above 500 and were considered positive for the presence of Strongylida type eggs. *Rhipicephalus sanguineus* ticks were collected from two animals. Sequences from the present study was placed together in a clade with *Mycoplasma* sp. and *Mycoplasma ovis* sequences from worldwide. Herein we report the occurrence of *Mycoplasma* sp. in sheep from Paraná State. Co-infection of gastrointestinal parasites and hemoplasmas may worse health condition of sheep, representing an alert about hemoplasmas diagnosis in ovine herds with nematode infections occurrence.

LIST OF FIGURES

PAPER FOR SUBMISSION

Figure 1: Phylogenetic tree based on partial sequences of the 16S rDNA gene, showing the relationship between the *Mycoplasma* sp. detected in sheep from this study and other hemoplasmas by Bayesian Inference. *Bacillus subtilis* and *Mycoplasma bovis* were used as outgroups. GenBank accession number is after the species name and origin of each bacterium. Analyses were carried out applying the F81+I+G model and 1000 bootstrap replicates for all analyses..... 49

APPENDIX

Figure 2: Phylogenetic tree based on partial 16S rRNA gene fragments of *Mycoplasma* sp. Tree was constructed by Bayesian Inference and a sequence from *Bacillus subtilis* was used as outgroup. Analysis were carried out applying the F81+I+G model and 1000 bootstrap replicates for all analyses. Our sequences are represented by ** after its name 61

LIST OF TABLES

OCCURRENCE OF HEMOPARASITES IN SHEEP

Table 1: Reports of hemoparasites molecularly detected in sheep	16
--	----

PAPER FOR SUBMISSION

Table 2: Obtained results regarding hematocrit values, amplification of the <i>Mycoplasma</i> spp. 16S rRNA fragment by PCR, EPG values and tick presence/identification in sheep from the present study.....	50
--	----

LIST OF ABBREVIATIONS

RBCs: Red blood cells

PCR: Polymerase chain reaction

PCV: Packed cell volume

GI: Gastrointestinal

EPG: Eggs per gram

B.C.: Before Christ

TBF: Tick-borne fever

DNA: Deoxyribonucleic acid

FEC: Fecal egg count

rDNA: Ribosomal deoxyribonucleic acid

TBD: Tick-borne diseases

mL: Milliliter

L/L: Liter of cells per liter of blood

EDTA: Ethylenediamine tetra acetic acid

GAPDH: Glyceraldehyde-3-phosphate dehydrogenase

bp: Base pair

BIC: Bayesian Information Criteria

CI: Confidence interval

NHP: Non-human primates

uL: Microliter

USA: United States of America

SUMMARY

1 INTRODUCTION	13
2 OBJECTIVES	15
2.1 General objectives	15
2.2 Specific objectives	15
3 CHAPTER ONE: OCCURRENCE OF HEMOPARASITES IN SHEEP	16
3.1 Sheep production.....	16
3.2. Hemoparasite infections in sheep.....	16
3.3 Anaplasmataceae	19
3.4 Piroplasmida	21
3.5 Hemotropic mycoplasmas	24
4 REFERENCES	25
5 CHAPTER TWO: MANUSCRIPT FOR SUBMISSION	38
5.1 Abstract	38
5.2. Introduction.....	38
5.3 Material and Methods.....	40
5.3.1 Ethical approval	40
5.3.2 Study	40
5.3.3 Sampling.....	40
5.3.4 Packed cell volume evaluation.....	41
5.3.5 Parasitological analysis	41
5.3.6 DNA extraction.....	41
5.3.7. Polymerase chain reactions (PCR)	41
5.3.8 Sequencing.....	42
5.3.9 Phylogenetic analysis	42
5.4 Results	42
5.6 Conclusion	45
5.7 References	45
6 APPENDIX: SECOND MANUSCRIPT FOR SUBMISSION	52
6.1 Introduction.....	52
6.2 Material and Methods.....	53
6.2.1 Sampling.....	53
6.2.2 DNA extraction and polymerase chain reaction (PCR) assays	54
6.2.3 Sequencing and phylogenetic analysis	54
6.3 Results	54
6.4 Discussion.....	55
6.5 Conclusion	57
6.6 References	57
7 FINAL CONCLUSIONS	63
REFERENCES	64

1 INTRODUCTION

Hemotropic mycoplasmas, also known as hemoplasmas, are small and pleomorphic bacteria of red blood cells (RBCs) that may infect many different species of vertebrate hosts (MESSICK et al., 2004). *Mycoplasma ovis* (formerly *Eperythrozoon ovis*) is a wall-less bacterium and it has been described infecting sheep (NEIMARK et al., 2004), goats (JOHNSON et al., 2016; MACHADO et al., 2017), wild ruminants (GRAZZIOTIN et al., 2011; GRAZZIOTIN et al., 2011b) and human beings (SYKES et al., 2010). In sheep, clinical manifestations of *M. ovis* infection may include weight loss (MARTINEZ-HERNANDEZ et al., 2018), hyperthermia, mucosal pallor and hemolytic anemia (NEIMARK et al., 2004; AKTAS, OZUBEK, 2017).

Diagnosis of sheep hemoplasmosis has been historically performed by visualization of compatible morphologic structures on RBCs during blood smears examination (NEITZ, 1934; SHERIFF et al., 1966; DADDOW, 1979; MASON and STATHAM, 1991). However, this diagnostic method may present low sensitivity and specificity when compared to molecular methods. Currently, molecular techniques, such as the polymerase chain reaction (PCR) with protocols aiming to detect genetic material from *M. ovis* has being used in sheep, goats and wild ruminants worldwide (OHTAKE et al., 2011; GRAZZIOTIN et al., 2011; GRAZZIOTIN et al., 2011b; BOES et al., 2012; HAMPEL et al., 2014; JOHNSON et al., 2016; AKTAS, OZUBEK, 2017; MACHADO et al., 2017; MARTINEZ-HERNANDEZ et al., 2018).

Mycoplasma ovis has been reported in sheep herds from different countries, such as Canada (NEIMARK et al., 2004); Japan (TAGAWA et al., 2012), Turkey (AKTAS, OZUBEK, 2017), China (WANG et al., 2017) and Hungary (HORNOK et al., 2018). In South America, only few reports of *M. ovis* infection in sheep from Argentina (AGUIRRE et al., 2009), and goats (MACHADO et al., 2017) and deer from Brazil (GRAZZIOTIN et al., 2011; GRAZZIOTIN et al., 2011b).

Brazilian sheep herd has been estimated in almost 18 million animals. Paraná State, located in the southern region of the country, has an estimated herd of more than 500,000 sheep (IBGE, 2017). Sheep farming has been considered a growing business for Paraná producers, most because of its profitability and demand (EMATER, 2018). Co-infection of *M. ovis* and other pathogens, such as nematodes and tick-borne agents, may lead hosts to develop anemia and may reflect on production decay and mortality in small ruminants

(HORNOK et al., 2009; ABDULLAH et al., 2013). Although the ovine production has a notable importance, studies regarding *M. ovis* occurrence in Brazilian sheep have not been performed to date. Thus, studies aiming to elucidate the distribution of *M. ovis* in Brazilian herds of sheep are needed.

2 OBJECTIVES

2.1 General objectives

To identify the causative agent of anemia in a sheep herd from Bandeirantes County, Paraná State, southern Brazil.

2.2 Specific objectives

- To collect and identify tick species parasitizing the animals;
- To determine the packed cell volume (PCV) on sheep samples;
- To screen sheep blood samples for *Mycoplasma* sp., *Anaplasma* sp. and piroplasm species by conventional PCR assays;
- To perform parasitological analysis of feces (worm egg counting – EPG);
- To molecularly characterize the pathogens detected by sequencing and comparison with other isolates deposited in the GenBank.
- To perform phylogenetic analysis with the isolates obtained in the present study aiming to elucidate its evolutionary features.

3 CHAPTER ONE: OCCURRENCE OF HEMOPARASITES IN SHEEP

3.1 Sheep production

Sheep, together with goats, are believed to be the first domesticated animals after dogs in approximately 9,000 years B.C. (ZYGOIYANNIS, 2006). Current domestic breeds were probably descendant from the wild mouflon (*Ovis aries orientalis*), originated from Asia (McMANUS et al., 2010). Worldwide herd is estimated in more than 1 billion animals, with the greater proportions concentrated in Asia, Oceania and Africa continents. South America represents 6.8% from the world sheep production (FAO, 2016). Although sheep are raised primarily for meat purposes, these animals provide a wide range of products, such as milky, skin and wool (ZYGOIYANNIS, 2006).

Regarding to Brazil, sheep were introduced by Portuguese colonizers resulting in well adapted breeds distributed through the country (PAIVA et al., 2005). Different sheep breeds may be found in Brazil, being all originated from exotic breeds or adapted in the country. Although these breeds present a variety of aptitudes, sheep production in Brazil is mainly directed to meat (MARIANTE et al., 2003).

In 2016, the Brazilian sheep herd reached the estimated number of almost 18,000,000 of animals added to an increasing long-term scenario of production and consumption of sheep products, foreseen by Brazilian researchers (EMBRAPA, 2017; IBGE, 2017). Paraná State, located in the southern region, have an estimated herd of more than 500,000 sheep, representing 3,12% from the national herd (IBGE, 2017) and considered a growing business (EMATER, 2018), counting with financial incentives from the government (FAEP, 2017).

3.2. Hemoparasite infections in sheep

A wide range of hemoparasites, such as bacteria and protozoa, are described infecting sheep from worldwide. Molecularly detected hemoparasite from sheep hosts found in the present review are summarized in Table 1.

Table 1. Reports of hemoparasites molecularly detected in sheep

Pathogen Group	Species	Occurrence	Reference
Anaplasmataceae	<i>Anaplasma phagocytophilum</i>	China, Czech Republic, France, Italy, Norway, Saudi Arabia, Slovakia.	Ladbury et al., 2008; Zhan et al., 2010; Torina et al., 2010; Derdakova et al., 2011; Dugat et al., 2014; Yang et al., 2015; Shabana et al., 2018.
	<i>A. phagocytophilum</i> -like	Tunisia	Ben Said et al., 2015; Ben Said et al., 2017.
	<i>A. ovis</i>	Algeria, China, Hungary, Iran, Iraq, Italy, Mongolia, Portugal, Saudi Arabia, Slovakia, South Africa, Sudan, Tunisia, Turkey.	Bekker et al., 2002; Hornok et al., 2007; Torina et al., 2010; Derdakova et al., 2011; Jalali et al., 2013; Renneker et al., 2013; Yang et al., 2015; Aouadi et al., 2017; Ochirkhuu et al., 2017; Yousefi et al., 2018; Aktas, Ozubek et al., 2018; Ringo et al., 2018; Shabana et al., 2018;
	<i>A. bovis</i>	China, Tunisia	Ben Said et al., 2015b; Zhang et al., 2016; Belkahia et al., 2017; Wang et al., 2018; Yang et al., 2018.
	<i>A. marginale</i>	Iran	Jalali et al., 2013; Yousefi et al., 2017.
	<i>A. platys</i> -like	Tunisia	Ghai et al., 2016; Ben Said et al., 2017b
	<i>A. capra</i>	China	Yang et al., 2018
	<i>Ehrlichia ruminantium</i>	Mozambique, South Africa	Bekker et al., 2002; Ringo et al., 2018.
Piroplasmids	<i>Theileria lestoquardi</i>	Sudan, Iran, Tunisia	Heidapour Bami et al., 2009; Zaeemi et al., 2011; Rjeibi et al., 2016b; Ali et al., 2017.
	<i>T. luwenshuni</i>	China, Turkey, Spain	Nagore et al., 2004; Yin et al., 2007; Chen et al., 2014; Ros-Garcia et al., 2016; Bilgic et al., 2017.

<i>T. uilenbergi</i>	China, Iran, Turkey	Yin et al., 2007; Renneker et al., 2013; Bilgic et al., 2017.
<i>T. ovis</i>	Algeria, China, Ethiopia, Pakista, Saudi Arabia, South Africa, Spain, Sudan, Tunisia, Turkey	Nagore et al., 2004; Altay et al., 2005; Zaeemi et al., 2011; Li et al., 2011; Shahzad et al., 2013; Gebrekidan et al., 2014; Auoadi et al., 2017; Ringo et al., 2018; Mghirbi et al., 2018; Alazani et al., 2018;
<i>Theileria</i> sp. OT3	Caribbean, China, Iran, Italy, Spain, Turkey	Nagore et al., 2004; Aydin et al., 2013; Tian et al., 2014; Giansgapero et al., 2015; Ros-Garcia et al., 2016; Zhang et al., 2015; Bilgic et al., 2017; Tabaei et al., 2018.
<i>T. separata</i>	Ethiopia, Sudan, Turkey	Gebrekidan et al., 2014; El Imam et al., 2016; Bilgic et al., 2017.
<i>Theileria</i> sp. MK	Turkey	Altay et al., 2012; Aydin et al., 2013; Ozubek, Aktas et al., 2016.
<i>T. annulata</i>	Iran, Turkey	Zaeemi et al., 2011; Ozubek, Aktas et al., 2016.
<i>Babesia ovis</i>	Iran, Iraq, Israel, Pakistan, Portugal, Spain, Tunisia, Turkey.	Aydin et al., 2013; Shahzad et al., 2013; Renneker et al., 2013b; Horta et al., 2014; Ros-Garcia et al., 2016; Erster et al., 2016; Rjeibi et al., 2016; Bazmani et al., 2018;
<i>B. motasi</i>	China, Italy, Spain	Nagore et al., 2004; Liu et al., 2007; Giangaspero et al., 2014; Ros-Garcia et al., 2016.

	<i>B. crassa</i>	Turkey, Iran	Schnittger et al., 2003; Bilgic et al., 2017.
	<i>B. crassa</i> -like	China	Jia et al., 2018.
	<i>B. lengau</i> -like	Greece	Giardini et al., 2012
Hemothropic mycoplasma	<i>Mycoplasma ovis</i>	Argentina, Canada, China, Hungary, Japan, Turkey	Hornok et al., 2004; Hornok et al., 2009; Aguirre et al., 2009; Tagawa et al., 2012; Aktas, Ozubek, 2017; Wang et al., 2017

3.3 Anaplasmataceae

Since 2001, Anaplasmataceae family is reorganized in four genera comprising *Anaplasma*, *Ehrlichia*, *Cowdria* and *Neorickettsia* species (DUMLER et al., 2001). Regarding this family, *Anaplasma phagocytophilum* and *Anaplasma ovis* are frequently described as disease causing in sheep (TORINA et al., 2010; RENNEKER et al., 2013). Sheep presenting poor health conditions are reported as more susceptible to be infected by multiple *Anaplasma* species (TORINA et al., 2010).

In 1932, a “tick-borne fever” (TBF) was described in sheep herds from the United Kingdom, causing febrile phases and loss of body weight in the animals. The infective agent was present in blood and spleen and its transmission was associated with *Ixodes ricinus* ticks (GORDON et al., 1932). The species *A. phagocytophilum*, a gram-negative obligatory bacterium, is responsible for infect granulocytes from a wide range of mammals and considered the causative agent of TBF in domestic ruminants (LALOY et al., 2009; RIKIHISA, 2011). Sheep may act as reservoirs for this pathogen in some areas, harboring persistent infections (THOMAS et al., 2012). These animals may play an important role in the epidemiological chain of *A. phagocytophilum*, being an infection source for susceptible ticks (OGDEN et al., 2002). Moreover, different genotypes may occur in the same flock (LADBURY et al., 2008; DUGAT et al., 2017) and wild animals may be involved in the epidemiological chain (ZHAN et al., 2010). Recently, intrauterine infection was also related to *A. phagocytophilum* persistently infected sheep (STUEN et al., 2018).

Although its occurrence in small ruminants is concentrated in European countries (LADBURY et al. 2008; WOLDEHIWET, 2010; STUEN, 2016), reports of *A. phagocytophilum* and *A. phagocytophilum*-like strains in sheep may be found in Asia (ZHAN et al., 2010; YANG et al., 2018), Middle East (SHABANA et al., 2018) and Africa (BEN SAID et al., 2015; BEN SAID et al., 2017). In Brazil, although some reports indicate genetically similar strains infecting wild animals (MONGRUEL et al., 2017), *A. phagocytophilum* occurrence was not effectively recognized.

Anaplasma ovis is a more selective pathogen, infecting mainly sheep and goats (KUTTLER, 1984) and it is considered a neglected infection of small ruminants that may lead to economic losses (RENNEKER et al., 2013). This pathogen infects erythrocytes and is often related to cause mild pathogenicity in sheep (HORNOK et al., 2007; RENNEKER et al., 2013). Nevertheless, cases of anemia may occur (YASINI et al., 2012) and animal condition may be aggravated when co-infection with other hemoparasites occur (AKTAS, OZUBEK, 2018). DNA from this pathogen have already been detected in milk from healthy sheep in China (ZHANG et al., 2016b). *Dermacentor andersoni* and *Rhipicephalus bursa* ticks are associated with transmission (KOCAN et al. 1991; DE LA FUENTE et al., 2005). *Rhipicephalus turanicus* tick may be also related to infections in sheep (AKTAS, OZUBEK, 2018).

Occurrence of *A. ovis* infecting sheep herds is reported from a different range of areas from the world, such as Algeria (AOUADI et al., 2017), Italy (DE LA FUENTE et al., 2005), Hungary (HORNOK et al., 2007), Iran (JALALI et al., 2013), Niger (DAHMANI et al., 2017), Turkey (AKTAS, OZUBEK, 2018) and China (YANG et al., 2015; ZHANG et al., 2016; YANG et al., 2018). In the United States, this pathogen is reported infecting wild ruminants (GOFF et al., 1993; DE LA FUENTE et al., 2006). *Anaplasma ovis* is considered highly involved in concurrent infections with other hemoparasites in sheep (JALALI et al., 2013; YANG et al., 2015; SEVINC et al., 2018; RINGO et al., 2018) and its occurrence is reported during all year seasons (BELKHIA et al., 2017). To date, there is no report of *A. ovis* occurring in any animal species in Brazil.

Other *Anaplasma* species are described infecting sheep around the world, as *Anaplasma capra*, a recently proposed new species that may also affect goats and humans in China (LI et al., 2015; YANG et al., 2018). Further investigations about this pathogen still needed. *Anaplasma marginale*, well recognized as one of the most important pathogens for cattle, occurs mostly in subtropical and tropical regions and infects cattle being transmitted by

Rhipicephalus microplus ticks (KOCAN et al., 2010). This pathogen was reported infecting sheep from Iran (JALALI et al., 2013; YOUSEFI et al., 2017) and, although there is no report of sheep infection in Brazil, goats were recently described being infected by *A. marginale* in the country (DA SILVA et al., 2018).

Strains related to *Anaplasma platys* were detected infecting sheep from Tunisia (GHAI et al., 2016; BEN SAID et al., 2017). This pathogen is known to infect platelets from dogs (DUMLER et al., 2001), although there are evidences of genetically close strains occurring in different host species, such as cats and buffaloes (ZOBBA et al., 2015; MACHADO et al., 2016). Infection by *A. platys*-like strains in small ruminants need further investigations. *Anaplasma bovis* is a monocyte parasite that mainly affects cattle, causing fever, anorexia, lymph node enlargement and mucosal pallor (MELO JUNIOR et al., 2010). However, its occurrence in sheep is frequently reported in China and Tunisia (BEN SAID et al., 2015b; ZHANG et al., 2016; BELKHIA et al., 2017; WANG et al., 2018; YANG et al., 2018.). In some studies, *A. bovis* prevalence in ovine showed to be even greater than *A. phagocytophilum* (ZHANG et al., 2016). A study conducted in Tunisia reported that *A. bovis* infection may be frequently mixed with *A. ovis* in small ruminants (BELKHIA et al., 2017).

Ehrlichia ruminantium, a gram-negative bacterium that replicates in endothelial cells, is considered the causative agent of heartwater. This pathogen affects domestic and wild ruminants mainly in the sub-Saharan Africa region (DUMLER et al., 2001; ALLSOPP et al., 2010). The disease name is given to pericardial effusions that may develop in small ruminants (ROBSON, ROBSON, 2016). Economic losses caused by the disease are considered expressive in affected areas (MELTEZER et al., 1996) and infections may lead mortalities rates above 90% in sheep (NEITZ et al., 1964). Its transmission is related to ticks from genus *Amblyomma* (BEKKER et al., 2002; ALLSOPP et al., 2010), but it was also found infecting *Rhipicephalus microplus* ticks (BIGUEZOTON et al., 2016). Mixed infections involving other blood parasites such as *A. ovis* and *Theileria ovis* may occur in sheep (RINGO et al., 2018). Although there are no reports about this agent infecting livestock in Brazil, a closely genotype of *Ehrlichia* sp. was found infecting a horse from the southern part of the country (VIEIRA et al., 2018).

3.4 Piroplasmida

Piroplasmids are protozoa agents belonging to order Piroplasmida. Regarding to ovine,

Babesia spp. and *Theileria* spp. represent the most important genera from this group with various species. These pathogens were considered neglected when compared to cattle piroplasmids. However, studies about small ruminants piroplasmids have increased together with the economic interest in these animals (SCHNITTGER et al., 2003).

Regarding ovine theileriosis, clinical signs in sheep may be presented as fever, cough, anorexia, mucosal pallor and ruminal hypomotility (HASSAN et al., 2015) and different species may act as the causative agent. Theileriosis transmissions in sheep are related to ticks from genera *Hyalomma* and *Rhipicephalus* (MORZARIA, 1998).

The species *Theileria lestoquardi*, considered the causative of the malignant theileriosis disease, is appointed as a highly pathogenic species, reaching morbidity and mortality rates of 100% and 90% in sheep, respectively (HOOSHMAND-RAD, HAWA, 1973). Its occurrence has been reported in Sudan, Iran and Tunisia (HEIDAPOUR-BAMI et al., 2009; ZAEEMI et al., 2011; RJEIBI et al., 2016b; ALI et al., 2017).

Phylogenetic analysis of *Theileria* sp. China 1 and *Theileria* sp. China 2 genotype revealed that these species genetically differ from *T. lestoquardi* and may present a pathogenic course in sheep. Moreover, a new classification of these species was proposed, being named as *Theileria luwenshuni* and *Theileria uilenbergi*, respectively (YIN et al., 2007). Once *Theileria* sp. OT1 genotype presented high genetical similarity with *Theileria* sp. China 1 it was also proposed as *T. luwenshuni* (ROS-GARCIA et al., 2013). Although *T. luwenshuni* and *T. uilenberg* were believed to circulate only in China, reports from other regions such as Spain (NAGORE et al., 2004), Iran (RENNEKER et al., 2013) and Turkey (BILGIC et al., 2017).

Some *Theileria* species appears to be less pathogenic in immunocompetent animals, such as *Theileria ovis* (ALANI, HEBERT, 1988; LI et al., 2011). Besides that, this species is frequently reported in different regions of the world, such as Spain (NAGORE et al., 2004), China (LI et al., 2011), Algeria (AOUADI et al., 2017) and South Africa (RINGO et al., 2018). Also, strain *Theileria* sp. OT3 have been isolated in sheep from Spain (NAGORE et al., 2004), Turkey (AYDIN et al., 2013), China (TIAN et al., 2014), Italy (GIANGASPERO et al., 2015), Caribbean Islands (ZHAG et al., 2015) and Iran (TABAEI et al., 2018). Although its widely occurrence, pathogenicity related to this strain is not fully elucidated but abortions may be associated (NAGORE et al., 2004). Tick species *R. sanguineus*, *Haemaphysalis punctata* and *Hyalomma detritum* are related as vectors (TIAN et al., 2014).

Species such as *Theileria separata*, *Theileria* sp. MK and *Theileria annulata* are scarcely reported in sheep by molecular techniques. *Theileria separata* is reported in sheep from Ethiopia (GEBREKIDAN et al., 2014), Sudan (EL IMAM et al., 2016) and Turkey (BILGIC et al., 2017), while *Theileria* sp. MK reports appears to be concentrated only in Turkey (ALTAY et al., 2012; AYDIN et al., 2013; OZUBEK, AKTAS et al., 2016). Although *T. annulata* is well recognized as a blood parasite from bovine hosts, it was reported occurring in sheep from Iran and Turkey (ZAEEMI et al., 2011; OZUBEK, AKTAS et al., 2016). Pathogenicity and vectors involved still need further investigations.

Although occurrence of *Theileria* species is described to occur in tropical and sub-tropical countries (HOOSHMAND-RAD, HAWA, 1973), reports appears to be concentrated in Africa (RJEIBI et al., 2016; ALI et al., 2017; RINGO et al., 2018), Asia (JIANXUN, HONG, 1997; ALTAY et al., 2005; YIN et al., 2007; AYDIN et al., 2013), Middle East (TAGELDIN et al., 2005; ZAEEMI et al., 2011; HASHEMINISAB et al., 2018) and some Europe regions (ALANI, HEBERT, 1988; GIANGASPERO et al., 2015). In Brazil, descriptions of *Theileria* detected in horses, buffaloes and wild animals (SILVEIRA et al., 2016; FERREIRA et al., 2016; SILVEIRA et al., 2017; VIEIRA et al., 2018b) were performed, but to date there are no reports about ovine theileriosis in the country.

Babesiosis is also caused by more than one species of *Babesia* spp. in sheep. Fever, haemoglobinuria and anemia are described as babesiosis clinical manifestations in ovine (YERUHAM, 1998). The disease is considered more life-threatening to sheep when compared to theileriosis and anaplasmosis (SEVINC et al., 2018). The species *Babesia ovis* and *Babesia motasi* are considered highly pathogenic for small ruminants (HAGHI et al., 2013) and transmitted by *Rhipicephalus* and *Haemaphysalis* ticks, respectively (UILENBERG, 2006). Besides that, vector species and pathogenicity of *Babesia crassa* still need further investigations (UILENBERG, 2006).

Species *B. ovis* is a wide distributed in Middle East, Europe and Asia (AYDIN et al., 2013; SHAHZAD et al., 2013; RENNEKER et al., 2013b; HORTA et al., 2014; ROS-GARCIA et al., 2013; ERSTER et al., 2016; RJEIBI et al., 2016; BAZMANI et al., 2018), meanwhile *B. motasi* and *B. crassa* are less reported using molecular tools. *Babesia crassa* is more related to reports from Asia and Middle East (SCHNITTGER et al., 2003; BILGIC et al., 2017) and *B. motasi* is described occurring in Asia and Europe (NAGORE et al., 2004; LIU et al., 2007; GIANGASPERO et al., 2014; ROS-GARCIA et al., 2013).

Furthermore, *Babesia* genotypes that remain not fully characterized are reported. A

genotype closely related to *Babesia lengau*, a novel species found in African cheetah (*Acinonyx jubatus*), are described infecting sheep from Greece that were presenting icteric membranes, anemia and slight leukocytosis (GIARDINI et al., 2012). *Babesia crassa*-like agents were identified in humans with clinical symptomatology in China, raising concerns about zoonotical transmission (JIA et al., 2018). In Brazil, although *Babesia* species are well documented in different animal hosts, there are no reports involving sheep until today.

3.5 Hemotropic mycoplasmas

The genus *Mycoplasma* is comprised within the Mollicutes class, and are distinguished by cell wall lackness (TRATCHENBERG, 2005). Species from *Haemobartonella* and *Eperythrozoon* genera were re-classified into the genus *Mycoplasma* after studies elucidating its real phylogenetic relationship (RIKIHISA et al., 1997; NEIMARK et al., 2001). Thus, classification of some *Mycoplasma* species as hemotropic mycoplasmas (hemoplasmas) were performed. These species, also known as hemoplasmas, are small and pleomorphic bacteria of red blood cells that may infect many different species of vertebrate hosts (MESSICK et al., 2004).

Since 2004, the recognized pathogen *Eperythrozoon ovis* has been described as *Mycoplasma ovis*. This species is described as a wall-less bacterium that infect sheep and goat erythrocytes (NEIMARK et al., 2004). Weight loss (MARTINEZ- HERNANDEZ et al., 2018), hyperthermia, mucosal pallor and hemolytic anemia (NEIMARK et al., 2004; AKTAS, OZUBEK, 2017) were clinical manifestations linked to *M. ovis* infection in ovine. Concurrent infections with internal parasites may increase the pathogenicity of *M. ovis* in sheep (OHTAKE et al., 2011).

Descriptions of *M. ovis* occurrence in sheep herds from different places of the world, such as Japan (TAGAWA et al., 2012), Turkey (AKTAS, OZUBEK, 2017), China (WANG et al., 2017) and Hungary (HORNOK et al., 2018) may be found. However, only few reports are originated from South America, being concentrated in Argentina (AGUIRRE et al., 2009). In Brazil, *M. ovis* has been reported in deer (GRAZZIOTIN et al., 2011; GRAZZIOTIN et al., 2011b) and goats (MACHADO et al., 2017). Sheep located in Japan that were originally from Australia were infected my hemotropic mycoplasmas (TAGAWA et al., 2012), raising a concern about sheep trade and pathogen distribution worldwide. Moreover, a *M. ovis*-like infecting a veterinarian has been found (SYKES et al., 2010), which

highlights for the zoonotical potential of this bacterium. Transmission route or the putative vector remains to be investigated.

4 REFERENCES

- ABDULLAH, J.; ADAMU, L.; JAMAL, M.H.A.B.; OSMAN, A.Y.; et al. Parasitic Gastro-Enteritis (PGE) Concurrent with Eperythrozoonosis in a Goat: A Case Report. **IOSR Journal of Agriculture and Veterinary Science**, v. 4, n. 1, p. 63-66, 2013.
- AGUIRRE, D. H.; THOMPSON, C.; NEUMANN, R. D.; SALATIN, A. O.; GAIDO, A. B. Brote de micoplasmosis clínica por *Mycoplasma ovis* en ovinos de Salta, Argentina. Diagnóstico clínico, microbiológico y molecular. **Revista Argentina de Microbiología**, v. 41, p. 212–214, 2009.
- AKTAS, M.; OZUBEK, S. *Anaplasma ovis* genetic diversity detected by major surface protein 1a and its prevalence in small ruminants. **Veterinary Microbiology**, v. 217, p. 13–17, 2018. AKTAS, M.; OZUBEK, S. Short communication A molecular survey of small ruminant hemotropic mycoplasmosis in Turkey, including first laboratory confirmed clinical cases caused by *Mycoplasma ovis*. **Veterinary Microbiology**, v. 208, p. 217–222, 2017.
- ALANI, A. J.; HERBERT, I. V. Pathogenesis of Infection with *Theileria recondita* (Wales) Isolated from *Haemaphysalis punctata* from North Wales. **Veterinary Parasitology**, v. 28, p. 293–301, 1988.
- ALAZANI, A.D.; SAID, E.A.; GHONEIM, A.M.; et al. A comprehensive evaluation and first molecular report of *Theileria ovis* infection in small ruminants in Saudi Arabia. **Tropical Animal Health and Production**, v. 51, n. 1, p. 89-98, 2018.
- ALI, A. M.; SALIH, D. A.; NJAHIRA, M. N.; et al. Genotyping of *Theileria lestoquardi* from sheep and goat in Sudan to support. **Veterinary Parasitology**, v. 239, p. 7–14, 2017.
- ALLSOPP, B. A. Natural history of *Ehrlichia ruminantium*. **Veterinary Parasitology**, v. 167, p. 123–135, 2010.
- ALTAY K., DUMANLI, N.; AKTAS, M. A study on ovine tick-borne hemoprotozoan parasites (*Theileria* and *Babesia*) in the East Black Sea Region of Turkey. **Parasitology Research**, v. 111, p. 149-153, 2012.
- ALTAY, K.; DUMANLI, N.; HOLMAN, P. J.; AKTAS, M. Detection of *Theileria ovis* in naturally infected sheep by nested PCR. **Veterinary Parasitology**, v. 127, p. 99–104, 2005.

- AOUADI, A.; LEULMI, H.; BOUCHEIKHCHOUKH, M.; BENAKHLA, A. Molecular evidence of tick-borne hemoprotozoan-parasites (*Theileria ovis* and *Babesia ovis*) and bacteria in ticks and blood from small ruminants in Northern Algeria. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 50, p. 34–39, 2017.
- AYDIN, M.F.; AKTAS, M.; DUMANLI, N. Molecular Identification of *Theileria* and *Babesia* in sheep and goats in the Black Sea Region in Turkey. **Parasitology Research**, v. 112, n. 8, p. 2817-2824, 2013.
- BAZMANI, A.; ABNOLHOOSHYAR, A.; IMANI-BARAN, A.; et al. Semi-nested polymerase chain reaction-based detection of *Babesia* spp. in small ruminants from Northwest of Iran. **Veterinary World**, v. 11, n. 3, p. 268-273, 2018.
- BEKKER, C. P. J.; VOS, S. DE; TAOU, A.; SPARAGANO, O. A. E.; JONGEJAN, F. Simultaneous detection of *Anaplasma* and *Ehrlichia* species in ruminants and detection of *Ehrlichia ruminantium* in *Amblyomma variegatum* ticks by reverse line blot hybridization. **Veterinary Microbiology**, v. 89, p. 223-238, 2002.
- BELKHIA, H.; BEN SAID, M.; MABROUK, N.E.; SAIDANI, M.; CHERNI, C.; HASSEN, M.B.; BOUATTOUT, A.; MESSADI, L. Seasonal dynamics, spatial distribution and genetic analysis of *Anaplasma* species infecting small ruminants from Northern Tunisia. **Infection, Genetics and Evolution**, v. 54, p. 66-73, 2017.
- BEN SAID, M.; BELKHIA, H.; ALBERTI, A.; et al. Molecular Survey of *Anaplasma* Species in Small Ruminants. **Vector-Borne and Zoonotic Diseases**, v. 15, n. 10, p. 1–11, 2015.
- BEN SAID, M.; BELKHIA, H.; KARAoud, M.; et al. First molecular survey of *Anaplasma bovis* in small ruminants from Tunisia. **Veterinary Microbiology**, v. 179, p. 322–326, 2015b.
- BEN SAID, M.; BELKHIA, H.; MABROUK, N. E.; et al. Molecular typing and diagnosis of *Anaplasma* spp. closely related to *Anaplasma phagocytophilum* in ruminants from Tunisia. **Ticks and Tick-borne Diseases**, v. 8, n. 3, p. 412–422, 2017.
- BIGUEZOTON, A.; NOEL, V.; ADEHAN, S.; et al. *Ehrlichia ruminantium* infects *Rhipicephalus microplus* in West Africa. *Parasites & Vectors*, n. 9, p. 4–7, 2016.
- BILGIC, H. B.; BAK, S.; KOSE, O.; et al. Prevalence of tick-borne haemoparasites in small ruminants in Turkey and diagnostic sensitivity of single-PCR and RLB. **Parasites & Vectors**, v. 10, p. 1–13, 2017.

- BOES, K. M.; GONCAROV, K. O.; THOMPSON, C. A.; et al. Identification of a *Mycoplasma ovis*-like organism in a herd of farmed white-tailed deer (*Odocoileus virginianus*) in rural Indiana. **Veterinary Clinical Pathology**, v. 41, n. 1, p. 77–83, 2012.
- CHEN, Z.; LIU, Q.; FENG-CHAO, J.; et al. Detection of piroplasms infection in sheep, dogs and hedgehogs in Central China. **Infectious Diseases of Poverty**, v. 3, p. 18–25, 2012.
- DA SILVA, N.B.; TAUS, N.S.; JOHNSON, W.C.; MIRA, A.; et al. First report of *Anaplasma marginale* infection in goats, Brazil. **PLoS One**, v. 13, n. 8, 2018.
- DADDOW, K.N. The transmission of a sheep strain of *Eperythrozoon ovis* to goats and the development of a carrier state in the goats. **Australian Veterinary Journal**, v. 55, p. 605, 1979.
- DAHMANI, M.; MARIÉ, J.; SCANDOLA, P.; BRAH, S. *Anaplasma ovis* infects sheep in Niger. **Small Ruminant Research**, v. 151, p. 32–35, 2017.
- DE LA FUENTE, J.; ATKINSON, M. W.; HOGG, J. T.; et al. Genetic Characterization of *Anaplasma ovis* Strains from Bighorn Sheep in Montana. **Journal of Wildlife Diseases**, v. 42, n. 2, p. 381–385, 2006.
- DE LA FUENTE, J.; TORINA, A.; CARACAPPA, S.; et al. Serologic and molecular characterization of *Anaplasma* species infection in farm animals and ticks from Sicily. **Veterinary Parasitology**, v. 133, p. 357–362, 2005.
- DERDAKOVA, M.; STEFANCIKOVA, A.; SPITALSKA, E.; et al. Emergence and genetic variability of *Anaplasma* species in small ruminants and ticks from Central Europe. **Veterinary Microbiology**, n. 153, p. 293–298, 2011.
- DUGAT, T.; CHASTAGNER, A.; LAGRÉE, A.; et al. A new multiple-locus variable-number tandem repeat analysis reveals different clusters for *Anaplasma phagocytophilum* circulating in domestic and wild ruminants. **Parasites & Vectors**, v. 7, p. 1–11, 2017.
- DUMLER, J. S.; BARBET, A. F.; BEKKER, C. P. J.; et al. Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designated of *Ehrlichia equi* and ‘HGE agent’ as subjective synonyms of *Ehrlichia phagocytophila*. **International Journal of Systematic Evolutionary Microbiology**, v. 51, n. 6, p. 2145–2165, 2001.
- EL IMAM, A.H.; HASSAN, S.M.; GAMEEL, A.A.; et al. Molecular identification of different *Theileria* and *Babesia* species infecting sheep in Sudan. **Annals of Parasitology**, v. 62, n. 2, p. 47–54, 2016.

EMATER. Instituto Paranaense de Assistência Técnica e Extensão Rural. Projetos Ovinos e Caprinos, 2018. Disponível online em: <http://www.emater.pr.gov.br/modules/conteudo/conteudo.php?conteudo=72>. (Acessado em 04 de Novembro de 2018).

EMBRAPA. Análise da PPM 2016: evolução dos rebanhos ovinos e caprinos entre 2007 e 2016. Boletim do Centro de Inteligência e Mercado de Caprinos e Ovinos, n. 1, p. 1-13, 2017. Disponível online em: <https://ainfo.cnptia.embrapa.br/digital/bitstream/item/165640/1/CNPC-2017-Boletim-CIM.n1.pdf>. (Acessado em 04 de Novembro de 2018).

ERSTER, O.; ROTH, A.; WOLLKOMIRSKY, R.; et al. Quantitative analysis of *Babesia ovis* infection in sheep and ticks. **Veterinary Parasitology**, v. 221, p. 39-45, 2016.

FAEP. Federação da Agricultura do Estado do Paraná. Incentivo aos Produtores de Caprinos e Ovinos, 2017. Disponível online em: <http://sistemafaep.org.br/incentivo-aos-produtores-de-ovinos-e-caprinos/>. (Acessado em 04 de Novembro de 2018).

FAO. Food and Agriculture Organization of the United Nations. FAOSTAT. Sheep. Live animals. Disponível online em: <http://www.fao.org/faostat/en/#search/sheep>. (Acessado em 10 de Dezembro de 2018).

FERREIRA, E. P.; VIDOTTO, O.; ALMEIDA, J. C.; et al. Serological and molecular detection of *Theileria equi* in sport horses of northeastern Brazil. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 47, p. 72-76, 2016.

GEBREKIDAN, H.; HAILU, A.; KASSAHUN, A.; ROHOU, I. *Theileria* infection in domestic ruminants in northern Ethiopia. **Veterinary Parasitology**, v. 200, p. 31-38, 2014.

GHAH, R.R.; MUTINDA, M.; EZENWA, V.O. Limited sharing of tick-borne hemoparasites between sympatric wild and domestic ungulates. **Veterinary Parasitology**, n. 226, p. 167-173, 2016.

GIANGASPERO, A.; MARANGI, M.; PAPINI, R.; et al. *Theileria* sp. OT3 and other tick-borne pathogens in sheep and ticks in Italy: Molecular characterization and phylogeny. **Ticks and Tick-borne Diseases**, v. 6, n. 1, p. 75-83, 2015.

GIARDINI, N.D.; CHOCHLAKIS, D.; KRITSEPI-KONSTANTINO, M.; et al. Haemolytic disease in sheep attributed to a *Babesia lengau*-like organism. **Veterinary Record**, v. 170, n. 6, p. 155-157, 2012.

GOFF, W.; STILLER, D.; JESSUP, D.; et al. Characterization of an *Anaplasma ovis* Isolate from Desert Bighorn Sheep in Southern California Desert Sheep Bighorn. **Journal of Wildlife Diseases**, v. 29, n. 4, p. 540-546, 1993.

GORDON, W. S.; BROWNLEE, A.; WILSON, D. R.; MACLEOD, J. Tick-Borne Fever: A

Hitherto Undescribed Disease of Sheep. **Journal of Comparative Pathology and Therapeutics**, v. 45, n. 2, p. 301–307, 1932.

GRAZZIOTIN, A. L.; DUARTE, J. M. B.; SZABÓ, M. P. J.; et al. Prevalence and Molecular Characterization of *Mycoplasma ovis* in Selected Free-Ranging Brazilian Deer Populations. **Journal of Wildlife Diseases**, v. 47, n. 4, p. 1005–1011, 2011.

GRAZZIOTIN, A. L.; SANTOS, A. P.; GUIMARAES, A. M. S.; et al. *Mycoplasma ovis* in captive cervids: Prevalence, molecular characterization and phylogeny. **Veterinary Microbiology**, v. 152, n. 3–4, p. 415–419, 2011b.

HAGHI, S.M.M.; FAKHAR, M.; SHARIF, M.; et al. Molecular Identification of Ovine Babesia sp. in North Iran. **Research in Molecular Medicine**, v. 1, n. 1, p. 35-39, 2013.

HAMPEL, J. A.; SPATH, S. N.; BERGIN, I. L.; et al. Prevalence and Diagnosis of Hemotrophic Mycoplasma Infection in Research Sheep and Its Effects on Hematology Variables and Erythrocyte Membrane Fragility. **Comparative Medicine**, v. 64, n. 6, p. 478–485, 2014.

HASHEMINISAB, S.S.; MORADI, P.; WRIGHT, I. A four-year epidemiological and chemotherapy survey of babesiosis and theileriosis, and tick vectors in sheep, cattle and goats in Dehgolan, Iran. **Annals of Parasitology**, v. 64, n. 1, p. 43-48, 2018.

HASSAN, M.A.; RAOOFI, A.; LOTFOLLAHZADEH, S.; JAVANBANKHT, J. Clinical and cytological characteristics and prognostic implications on sheep and goat *Theileria* infection in north of Iran. **Journal of Parasitic Diseases**, v. 39, n. 2, p. 190-193, 2015.

HEIDAPOUR-BAMI, M.; HADDADZADEH, H.R.; KAZEMI, B.; et al. Molecular identification of ovine *Theileria* species by a new PCR-RFLP method. **Veterinary Parasitology**, v. 12, n. 3-4, p. 171-177, 2009.

HOOSHMAND-RAD, P.; HAWA, N. J. Malignant Theileriosis of Sheep. **Tropical Animal Health and Production**, v. 5, p. 97–102, 1973.

HORNOK, N.; HOFF, B.; GANTER, M. *Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an eperythrocyclic agent of haemolytic anemia in sheep and goats. **International Journal of Systematic and Evolutionary Microbiology**, v. 54, p. 365-71, 2004.

HORNOK, S.; ELEK, V.; DE LA FUENTE, J.; NARANJO, V.; FARKAS, R.; MAJOROS, G.; FOLDVÁRI, G. First serological and molecular evidence of endemicity of *Anaplasma ovis* and *A. marginale* in Hungary. **Veterinary Microbiology**, v. 122, p. 316-322, 2007.

HORNOK, S.; MELI, M. L.; ERDOS, A.; et al. Molecular characterization of two different

- strains of haemotropic mycoplasmas from a sheep flock with fatal haemolytic anaemia and concomitant *Anaplasma ovis* infection. **Veterinary Microbiology**, v. 136, p. 372–377, 2009.
- HORNOK, S.; SUGAR, L.; FERNÁNDEZ, I.F.; et al. Tick- and fly-borne bacteria in ungulates: the prevalence of *Anaplasma phagocytophilum*, haemoplasmas and rickettsiae in water buffalo and deer species in Central Europe, Hungary. **BMC Veterinary Research**, v. 14, p. 98-104, 2018.
- HORTA, S.; BARRETO, M.C.; PEPE, A.; CAMPOS, J.; OLIVA, A. Highly sensitivity method for diagnosis of subclinical *B. ovis* infection. **Ticks and Tick-Borne Diseases**, v. 5, p. 902-906, 2014.
- IBGE. Pesquisa da Pecuária Municipal. Efetivos de rebanho, por tipo de rebanho, 2017. Disponível online em: <https://sidra.ibge.gov.br/tabela/3939>. (Acessado em 22 de Julho de 2018).
- JALALI, R. Molecular detection and identification of *Anaplasma* species in sheep from Ahvaz, Iran. **Iranian Journal of Veterinary Research**, v. 14, n. 1, p. 50–56, 2013.
- JIA, N.; ZHENG, Y.; JIANG, J.; et al. Human babesiosis caused by a novel *Babesia crassa*-like pathogen: A case series. **Clinical Infectious Diseases**, v. 67, n. 7, p. 110–1119, 2018.
- JIANXUN, L; HONG, Y. Theileriosis of Sheep and Goats in China. **Tropical Animal Health Production**, v. 29, p. 85-105, 1997.
- JOHNSON, K. A.; BAUER, A. E.; WENG, H.; et al. Detection of hemoplasma infection of goats by use of a quantitative polymerase chain reaction assay and risk factor analysis for infection. **American Journal of Veterinary Research**, v. 77, n. 8, 2016.
- KOCAN, K. M. et al. The natural history of *Anaplasma marginale*. **Veterinary Parasitology**, v. 167, n. 2-4, p. 95-107, 2010.
- KOCAN, K.M.; STILLER, D. Development of *Anaplasma ovis* (Rickettsiales: Anaplasmataceae) in male *Dermacentor andersoni* (Acari: Ixodidae) transferred from infected to susceptible sheep. **Journal of Medical Entomology**, v. 29, n. 1, p. 98-107, 1991.
- KUTTLER, A. K. L. *Anaplasma* Infections in Wild and Domestic Ruminants: A Review. **Journal of Wildlife Diseases**, v. 20, n. 1, p. 12–20, 1984.
- LADBURY, G. A. F.; STUEN, S.; THOMAS, R.; et al. Dynamic Transmission of Numerous *Anaplasma phagocytophilum* Genotypes among Lambs in an Infected Sheep Flock in an Area of Anaplasmosis Endemicity. **Journal of Clinical Microbiology**, v. 46, n. 5, p. 1686–1691, 2008.

- LALOY, E.; PETIT, E.; BOULOUIS, H.; et al. Dynamics of natural infection by *Anaplasma phagocytophilum* in a dairy cattle herd in Brittany, France. **Clinical Microbiology and Infection**, v. 15, p. 24–25, 2009.
- LI, H.; ZHENG, Y.; MA, L.; et al. Human infection with a novel tick-borne *Anaplasma* species in China: a surveillance study. **The Lancet Infectious Diseases**, v. 15, n. 6, p. 663–670, 2015.
- LI, Y.; GUAN, G.; MA, M.; LIU, J.; REN, Q.; LUO, J.; YIN, H. *Theileria ovis* discovered in China. **Experimental Parasitology**, n. 127, p. 304–307, 2011.
- LIU, A. H.; YIN, H.; GUAN, G. Q.; et al. At least two genetically distinct large *Babesia* species infective to sheep and goats in China. **Veterinary Parasitology**, v. 147, p. 246–251, 2007.
- MACHADO, C. A. L.; VIDOTTO, O.; CONRADO, F. O.; et al. *Mycoplasma ovis* infection in goat farms from northeastern Brazil. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 55, n. July, p. 1–5, 2017.
- MACHADO, R. Z.; MARIA, M.; TEIXEIRA, G.; et al. Molecular diagnosis and genetic diversity of tick-borne Anaplasmatidae agents infecting the African buffalo *Syncerus caffer* from Marromeu Reserve in Mozambique. **Parasites & Vectors**, v. 9, p. 454–463, 2016.
- MARIANTE, A.S.; MCMANUS, C.; MENDONÇA, J.F. **Country Report on the State of Animal Genetics Resources Brazil**. 1^o Edição. Brasília: Embrapa Genetic Resources and Biotechnology, 2003.
- MARTINEZ-HERNÁNDEZ, J. M.; BALLADOS-GONZÁLEZ, G. G.; FERNÁNDEZ-BANDALA, D.; et al. Molecular detection of *Mycoplasma ovis* in an outbreak of hemolytic anemia in sheep from Veracruz, Mexico. **Tropical Animal Health and Production**, n. 51, n. 4, p. 243–248, 2019.
- MASON, R.W.; STATHAM, P. Susceptibility of sheep and goats to *Eperythrozoon ovis* infection. **Australian Veterinary Journal**, v. 68, p. 116–117, 1991.
- MCMANUS, C.; PAIVA, S. R.; ARAÚJO, R. O. DE. Genetics and breeding of sheep in Brazil Genética e melhoramento de ovinos no Brasil. **Revista Brasileira de Zootecnia**, v.39, p. 236– 246, 2010.
- MELO-JUNIOR, O.A.; ALBERNAZ, A.P.; THIEBAUT, J.T.L. et al. Ocorrência de *Anaplasma bovis* (Donatien & Lestoquard, 1936, Dumler et al., 2001) na região de Bom Jesus do Itabapoana, RJ. **Ciência Animal Brasileira**, v. 11, n. 4, p.888–8891, 2010.
- MELTEZER, M.I.; PERRY, B.D.; DONACHIE, P.L. Mortality percentages related to

- heartwater and the economic impact of heartwater disease on large-scale commercial farms in Zimbabwe. **Preventive Veterinary Medicine**, v. 26, p. 187-199, 1996.
- MESSICK, J. B. Hemotrophic mycoplasmas (hemoplasmas): a review and new insights into pathogenic potential. **Veterinary Clinical Pathology**, v. 33, n. 1, p. 20–22, 2004.
- TABAEI, S.J.S.; SPOTIN, A.; POURIRAN, R.; et al. Molecular characterization of *Theileria* spp. in livestock and the first report on the occurrence of *Theileria* sp. OT3 in Iran. **Acta Parasitologica**, v. 63, n. 3, p. 515–521, 2018.
- MGHIRBI, Y.; ROS-GARCIA, A.; IRIBAR, P.; et al. A molecular study of tick-borne haemoprotozoan parasites (*Theileria* and *Babesia*) in small ruminants in Northern Tunisia. **Veterinary Parasitology**, v. 198, p. 72-77, 2013.
- MONGRUEL, A. C. B.; BENEVENUTE, J. L.; IKEDA, P.; et al. Detection of *Anaplasma* sp. phylogenetically related to *A. phagocytophilum* in a free-living bird in Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 26, n. 4, p. 505–510, 2017.
- MORZARIA, S.P. Theileriosis. In: DELVES, P.J; ROITT, I. **Encyclopedia of Immunology**. São Diego, CA, Estados Unidos: Academic Press, 1998.
- NAGORE, D.; GARCÍA-SANMARTÍN, J.; GARCÍA-PÉREZ, A. L.; et al. Identification, genetic diversity and prevalence of *Theileria* and *Babesia* species in a sheep population from Northern Spain. **International Journal for Parasitology**, v. 34, p. 1059–1067, 2004.
- NEIMARK, H.; HOFF, B.; GANTER, M. *Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an eperythrocytic agent of haemolytic anaemia in sheep and goats. **International Journal of Systematic and Evolutionary Microbiology**, v. 54, p. 365–371, 2004.
- NEIMARK, H.; JOHANSSON, K.; RIKIHISA, Y.; et al. Proposal to transfer some members of the genera *Haemobartonella* and *Eperythrozoon* to the genus *Mycoplasma* with descriptions of ‘*Candidatus Mycoplasma haemofelis*’, ‘*Candidatus Mycoplasma haemomuris*’, ‘*Candidatus Mycoplasma haemosuis*’ and ‘*Candidatus*. **International Journal of Systematic and Evolutionary Microbiology**, v. 51, p. 891–899, 2001.
- NEITZ, W. O.; ALEXANDER, R. A.; DU TOIT, P. J. *Eperythrozoon ovis* (sp. nov.) Infection in Sheep. **Onderstepoort Journal of Veterinary Science and Animal Industry**, v. 3, n. 2, p. 263–271, 1934.
- NEITZ, W.O. Tick-borne diseases as a hazard in the rearing of calves in Africa. **Bulletin de l’Office International des Épidémiologies**, v. 62, p. 607-625, 1964.

- OCHIRKHUU, N.; KONNAI, S.; ODBILEG, R.; et al. Molecular Epidemiology Survey and Genetic Characterization of *Anaplasma* Species in Mongolian Livestock. **Vector Borne and Zoonotic Diseases**, n. 17, v. 8, p. 539-549, 2017.
- OGDEN, N. H.; CASEY, A. N. J.; FRENCH, N. P.; et al. Natural *Ehrlichia phagocytophila* transmission coefficients from sheep ‘carriers’ to *Ixodes ricinus* ticks vary with the numbers of feeding ticks. **Parasitology**, v. 124, p. 127–136, 2002.
- OHTAKE, Y.; NISHIZAWA, I.; SATO, M.; WATANABE, Y.; NISHIMURA, T. *Mycoplasma ovis* Detected in Free-Living Japanese Serows, *Capricornis crispus*. **Journal of Veterinary Medicine and Science**, v. 73, n. 3, p. 371–373, 2011.
- OZUBEK, S.; AKTAS, M. Molecular and Parasitological Survey of Ovine Piroplasmosis, Including the First Report of *Theileria annulata* (Apicomplexa: Theileridae) in Sheep and Goats from Turkey. **Journal of Medical Entomology Advance**, v. 0, n. 0, p.1-9, 2016.
- PAIVA, D. F.; SILVÉRIO, V.C.; PAIVA, D.A.F.; MCMANUS, C.; et al. Origin of the Main Locally Adapted Sheep Breeds of Brazil: A RFLP-PCR Molecular Analysis. **Archivos de Zootecnia**, v. 54, p. 395-399, 2005.
- RENNEKER, S.; ABDO, J.; BAKHEIT, M.; et al. Coinfection of Sheep with *Anaplasma*, *Theileria* and *Babesia* Species in the Kurdistan Region, Iraq. **Transboundary and Emerging Diseases**, v. 60, b. 2, p. 113-118, 2013b.
- RENNEKER, S.; ABDO, J.; SALIH, D. E. A.; KARAGENC, T. Can *Anaplasma ovis* in Small Ruminants be Neglected any Longer? **Transboundary and Emerging Diseases**, v. 60, p. 105– 112, 2013.
- RIKIHISA, Y. Mechanisms of Obligatory Intracellular Infection with *Anaplasma phagocytophilum*. **Clinical Microbiology Reviews**, v. 24, n. 3, p. 469–489, 2011.
- RIKIHISA, Y.; KAWAHARA, M.; WEN, B.; et al. Western Immunoblot Analysis of *Haemobartonella muris* and Comparison of 16S rRNA Gene Sequences of *H. muris*, *H. felis*, and *Eperythrozoon suis*. **Journal of Clinical Microbiology**, v. 35, n. 4, p. 823–829, 1997.
- RINGO, A. E.; MOUMOUNI, P. F. A.; TAIÖE, M.; et al. Molecular analysis of tick-borne protozoan and rickettsial pathogens in small ruminants from two South African provinces. **Parasitology International**, v. 67, n. 2, p. 144–149, 2018.
- RJEIBI, M. R.; DARGHOUTH, M. A.; REKIK, M.; et al. First Molecular Identification and Genetic Characterization of *Theileria lestoquardi* in Sheep of the Maghreb Region. **Transboundary and Emerging Diseases**, v. 63, p. 278–284, 2016b.
- RJEIBI, M. R.; DARGHOUTH, M.; GHARBI, M. Prevalence of *Theileria* and *Babesia*

species in Tunisian sheep. **Onderstepoort Journal of Veterinary Research**, v. 83, n. 1, p. a1040, 2016.

ROBINSON, W.F.; ROBINSON, N.A. Chapter 1 – Cardiovascular System. In: JUBB, K.V.F.; KENNEDY, P.C.; PALMER, N. **Pathology of domestic animals**. São Diego, CA, Estados Unidos: Academic Press, 1993. p. 1-101.

ROS-GARCIA, A.; BARANDIKA, J.F.; GARCÍA-PEREZ, A.L.; JUSTE, R.A.; HURTADO, A. Assessment of exposure to piroplasms in sheep grazing in communal mountain pastures by using a multiplex DNA bead-based suspension array. **Parasites & Vectors**, v. 6, p. 277-288, 2013.

SCHNITTGER, L.; YIN, H.; GUBBELS, M. J.; et al. Phylogeny of sheep and goat *Theileria* and *Babesia* parasites. **Parasitology Research**, v. 91, p. 398–406, 2003.

SEVINC, F.; ZHOU, M.; CAO, S.; et al. Haemoparasitic agents associated with ovine babesiosis: A possible negative interaction between *Babesia ovis* and *Theileria ovis*. **Veterinary Parasitology**, v. 252, n. January, p. 143–147, 2018.

SHABANA, I. I.; ALHADLAG, N. M.; ZARAKET, H. Diagnostic tools of caprine and ovine anaplasmosis: a direct comparative study. **BMC Veterinary Research**, v.14, p. 1–8, 2018.

SHAHZAD, W.; NOOR, H.; AHMAD, M.; et al. Prevalence and Molecular Diagnosis of *Babesia ovis* and *Theileria ovis* in Lori Sheep at Livestock Experiment Station (LES), Bahadurnagar, Okara, Pakistan. **Iranian Journal of Parasitology**, v. 8, n. 4, p. 570-578, 2013.

SHERIFF, D.; CLAPP, K. H.; REID, M. *Eperythrozoon ovis* INFECTION IN SOUTH AUSTRALIA. **Australian Veterinary Journal**, v. 42, p. 169, 1966.

SILVEIRA, A. W.; OLIVEIRA, G. G.; SANTOS, L. M.; et al. Natural infection of the South America Tapir (*Tapirus terrestris*) by *Theileria equi*. **Journal of Wildlife Diseases**, v. 53, n. 2, p. 411–413, 2017.

SILVEIRA, J. A. G.; OLIVEIRA, C. H. S. DE; SILVESTRE, B. T.; et al. Molecular assays reveal the presence of *Theileria* spp. and *Babesia* spp. in Asian water buffaloes (*Bubalus bubalis*, Linnaeus, 1758) in the Amazon region of Brazil. **Ticks and Tick-borne Diseases**, v. 7, n. 5, p. 1017–1023, 2016.

STUEN, S. Tick-Borne Fever (*Anaplasma phagocytophilum* Infection) in Sheep – A Review. **Journal of Veterinary Medicine and Research**, v. 3, 2016.

- STUEN, S.; OKSTAD, W.; SAGEN, A. M. Intrauterine Transmission of *Anaplasma phagocytophilum* in Persistently Infected Lambs. **Veterinary Sciences**, v. 5, p.25-32, 2018.
- SYKES, J.E.; LINDSAY, L.L.; MAGGI, R.G.; BREITSCHWERDT, E. Human Coinfection with *Bartonella henselae* and Two Hemotropic Mycoplasma Variants Resembling *Mycoplasma ovis*. **Journal of Clinical Microbiology**, v. 48, n. 10, p. 3782-3785, 2010.
- TAGAWA, M.; TAKEUCHI, T.; FUJISAWA, T.; et al. A Clinical Case of Severe Anemia in a Sheep Coinfected with *Mycoplasma ovis* and ‘*Candidatus Mycoplasma haemovis*’ in Hokkaido, Japan. **Journal of Veterinary Medicine and Science**, v. 74, n. 1, p. 99-102, 2012.
- TAGELDIN, M. H.; FADIYA, A. A.; SABRA, A. A.; ISMAILY, S. I. A. Theileriosis in sheep and goats in the Sultanate of Oman. **Tropical Animal Health and Production**, v. 37, p. 491– 493, 2005.
- THOMAS, R. J.; BIRTLES, R. J.; RADFORD, A. D.; WOLDEHIWET, Z. Recurrent Bacteraemia in Sheep Infected Persistently with *Anaplasma phagocytophilum*. **Journal of Comparative Pathology**, v. 147, n. 2–3, p. 360–367, 2012.
- TIAN, Z.; LIU, G.; YIN, H.; et al. Parasitology International First report on the occurrence of *Theileria* sp. OT3 in China. **Parasitology International**, v. 63, n. 2, p. 403–407, 2014.
- TORINA, A.; GALINDO, R. C.; VICENTE, J.; et al. Characterization of *Anaplasma phagocytophilum* and *A. ovis* infection in a naturally infected sheep flock with poor health condition. **Tropical Animal Health and Production**, v. 42, p. 1327–1331, 2010.
- TRATCHENBERG, S. Quick Guide: Mollicutes. **Current Biology**, v. 15, n. 12, p. PR483-484, 2005.
- UILENBERG, G. *Babesia* - A historical overview. **Veterinary Parasitology**, v. 138, p. 3–10, 2006.
- VIERIA, M. I. B.; COSTA, M. M.; OLIVEIRA, M. T.; et al. Serological detection and molecular characterization of piroplasmids in equids in Brazil. **Acta Tropica**, v. 179, p. 81-87, 2018b.
- VIEIRA, T. S. W. L.; QUROLLO, B. A.; MONGRUEL, A. C. B.; et a. Potentially Same Novel *Ehrlichia* Species in Horses in Nicaragua and Brazil. **Emerging Infectious Diseases**, v. 24, n. 5, p. 180238, 2018.

- WANG, J.; ZHANG, Y.; CUI, Y. et al. A rapid, simple and sensitive loop-mediated isothermal amplification method to detect *Anaplasma bovis* in sheep and goat samples. **Parasitology International**, n. 67, p. 70-73, 2018.
- WANG, X.; CUI, Y.; ZHANG, Y.; et al. Molecular characterization of hemotropic mycoplasmas (*Mycoplasma ovis* and ‘*Candidatus Mycoplasma haemovis*’) in sheep and goats in China. **BMC Veterinary Research**, v. 13, p. 142-150, 2017.
- WOLDEHIWET, Z. The natural history of *Anaplasma phagocytophilum*. **Veterinary Parasitology**, v. 167, p. 108-122, 2010.
- YANG, J.; HAN, R.; NIU, Q.; et al. Ticks and Tick-borne Diseases Occurrence of four *Anaplasma* species with veterinary and public health significance in sheep, northwestern China. **Ticks and Tick-borne Diseases**, v. 9, n. 1, p. 82–85, 2018.
- YANG, J.; LI, Y.; LIU, Z.; et al. Molecular detection and characterization of *Anaplasma* spp. in sheep and cattle from Xinjiang, northwest China. **Parasites & Vectors**, v. 8, p. 108–115, 2015.
- YASINI, S.; KHAKI, Z.; RAHBARI, S.; KAZEMI, B. Hematologic and Clinical Aspects of Experimental Ovine Anaplasmosis Caused by *Anaplasma ovis* in Iran. **Iranian Journal of Parasitology**, v. 7, n. 4, p. 91-98, 2012.
- YERUHAM, I.; HADANI, A.; GALKER, F. Some epizootiological and clinical aspects of ovine babesiosis caused by *Babesia ovis* — a review. **Veterinary Parasitology**, v. 74, p. 153- 163, 1998.
- YIN, H.; SCHNITTGER, L.; LUO, J.; SEITZER, U.; AHMED, J. S. Ovine theileriosis in China: a new look at an old story. **Parasitology Research**, v. 101, n. 2, p. 191–195, 2007.
- YOUSEFI, A.; RAHBARI, S.; SHAYAN, P. et al. Molecular detection of *Anaplasma marginale* and *Anaplasma ovis* in sheep and goat in west highland pasture of Iran. **Asian Pacific Journal of Tropical Biomedicine**, v. 7, n. 5, p. 455- 459, 2017.
- ZAEEMI, M.; HADDADZADEH, H. Identification of different *Theileria* species (*Theileria lestoquardi*, *Theileria ovis*, and *Theileria annulata*) in naturally infected sheep using nested PCR – RFLP. **Parasitology Research**, v. 108, p. 837–843, 2011.
- ZHAN, L.; CAO, W.; JIANG, J.; et al. *Anaplasma phagocytophilum* from Rodents and Sheep, China. **Emerging Infectious Diseases**, v. 16, n. 5, 2010.
- ZHANG, J.; KELLY, P.; LI, J.; et al. Molecular detection of *Theileria* spp. in Livestock in Caribbean Islands. **BioMed Research International**, v. 8, p. 1-8, 2015.
- ZHANG, Y.; LV, Y.; CUI, Y.; et al. First molecular evidence for the presence of *Anaplasma*

DNA in milk from sheep and goats in China. **Parasitology Research**, v. 115, n. 7, p. 1–7, 2016b.

ZHANG, Y.; LV, Y.; ZHANG, F.; ZHANG, W. Molecular and phylogenetic analysis of *Anaplasma* spp. in sheep and goats from six provinces of China. **Journal of Veterinary Science**, v. 17, n. 4, p. 523-529, 2016.

ZOBBA, R.; ANFOSSI, A. G.; VISCO, S.; et al. Ticks and Tick-borne Diseases Cell tropism and molecular epidemiology of *Anaplasma platys* -like strains in cats. **Ticks and Tick-borne Diseases**, v. 6, n. 3, p. 272–280, 2015.

ZYGOYIANNIS, D. Sheep production in the world and in Greece. **Small Ruminant Research**, v. 62, p. 143–147, 2006.

5 CHAPTER TWO: MANUSCRIPT FOR SUBMISSION

Title: Haplotype diversity of *Mycoplasma ovis* in sheep from Paraná state, southern Brazil.

5.1 Abstract

Mycoplasma ovis is a wall-less bacterium that infects sheep and goat erythrocytes. Infection by *M. ovis* in sheep may lead to weight loss, submandibular edema, hyperthermia, mucosal pallor and hemolytic anemia. Although the importance of sheep industry in Brazil, no studies regarding *M. ovis* infection have been reported in the country. Moreover, *Anaplasma marginale* has been detected in goats from Brazil. Accordingly, the aims of this study were to i) screen sheep blood samples for *Mycoplasma* sp. and the tick-borne agents *A. marginale* and piroplasms infection by PCR; ii) identify the tick species parasitizing the animals; iii) evaluate the presence of co-infection with gastrointestinal (GI) nematodes by McMaster Fecal Egg Counting (FEC); and iv) determine the association between the presence of anemia and infection by these agents in a sheep herd from Bandeirantes City, Paraná State, southern Brazil. Seven out of 42 (16.6%; 95% CI: 8.32–30.6%) sheep were positive for *Mycoplasma* spp. and all tested negative for *A. marginale* and piroplasms by PCR. Two animals were infested by ticks. FEC was performed in 38 sheep and 24 (63.15%; 95% CI: 47.2 – 76.6%) presented > 500 eggs per gram (EPG). Nine out of 42 (21.42%; 95%CI: 11.71-35.94%) animals were anemic. Sequencing of three *Mycoplasma*-positive sample showed $\geq 99\%$ identity to multiple *M. ovis* 16S rDNA gene sequences deposited in GenBank. Obtained sequences clustered together with *Mycoplasma* sp. and *Mycoplasma ovis* through Bayesian inference. Haplotype analysis showed three different genotypes. *Mycoplasma* sp. occurs in sheep from the north region of Paraná State and may develop anemia in co-infected animals.

Key-words: Small ruminants, hemoplasmas, *Mycoplasma ovis*, *Anaplasma marginale*, piroplasms.

5.2. Introduction

Brazilian sheep herd has been estimated in more than 18 million animals. The southern region of the country concentrates 30% of the Brazilian herd, which are mainly focused on

meat production (IBGE, 2016). The gastrointestinal (GI) parasite *Haemonchus contortus* is responsible for great impact to the sheep industry (Lane et al., 2015), since it infects the sheep abomasum, causing an acute hemorrhagic anemia crisis which may leads to sudden death (Taylor et al., 2007). In sheep, although anemia is commonly linked to *H. contortus* infection, it is important to state that other infectious agents, such as hemotropic mycoplasmas (hemoplasmas) and tick-borne diseases (TBDs) agents, may also leads to anemic crisis (Yeruham et al., 1998; Neimark et al., 2004; Hornok et al., 2009; Alessandra and Santo, 2012), and historically have not been included as differential diagnosis of anemia by veterinary practitioners in Brazil.

Hemoplasmas are small and pleomorphic bacteria of red blood cells that may infect many different species of vertebrate hosts (Messick et al., 2004). In small ruminants, two hemoplasma species have been initially found *Mycoplasma ovis* (formerly *Eperythrozoon ovis*) and ‘*Candidatus Mycoplasma haemovis*’ (Hornok et al., 2012). However, the complete genome sequence of *M. ovis* strain Michigan revealed two copies of the 16S rDNA genes, which corresponded to the previously reported sequences for *M. ovis* and ‘*Ca. M. haemovis*’ (Deshuillers et al., 2014). In sheep, *M. ovis* infection may cause weight loss, hyperthermia, mucosal pallor and hemolytic anemia (Neimark et al., 2004; Aktas and Ozubek, 2017; Martinez-Hernandez et al., 2019). In Brazil, *M. ovis* has been detected in deer (Grazziotin et al., 2011a; 2011b) and goats (Machado et al., 2017).

Rhipicephalus microplus ticks, are endemic in Brazil (Dantas-Torres et al., 2009). Despite host specificity of *R. microplus* for cattle (Ma et al., 2016), this tick species may be found parasitizing small ruminants (Brito et al., 2005), and thus, may transmit pathogens. In Brazil, *Anaplasma marginale*, the causative agent of bovine anaplasmosis, is transmitted by *R. microplus* ticks and this bacterium has been detected in goats from the northeastern region of the country (Da Silva et al., 2018). Additionally, this bacterium has been detected in sheep from Iran (Yousefi et al., 2017). However, the epidemiological and clinical impact of *A. marginale* infection in small ruminants remains to be fully established.

A wide range of piroplasm species, mainly from the genus *Theileria* and *Babesia*, may infect sheep. Although piroplasmid infections in small ruminants may be considered neglected when compared to cattle infections, studies have increased together with the economic interest in these animals (Schnittger et al., 2003). These infections may cause fever, anorexia, mucosal pallor, hemoglobinuria and anemia (Yeruham et al., 1998; Hassan et al., 2015). *Rhipicephalus*, *Hyalomma* and *Haemaphysalis* ticks are related as putative vectors (Morzaria, 1998; Tian et

al., 2004; Uilenberg, 2006). In Brazil, studies on piroplasmid infection in sheep have not been reported.

Co-infection between *M. ovis* and other pathogens, such as nematodes and TBDs agents, may leads to hemolytic anemia and reflect on production decay and mortality (Hornok et al., 2009; Abdullah et al., 2013). Although the ovine production has a notable importance, no studies on *M. ovis*, *A. marginale* and piroplasms occurrence in sheep from Brazil have been reported. Accordingly, the aims of this study were to i) screen a sheep herd for *Mycoplasma* sp. and the TBDs agents *A. marginale* and piroplasms by PCR; ii) identify the tick species parasitizing the animals; iii) evaluate the presence of co-infection with gastrointestinal (GI) nematodes by McMaster Fecal Egg Counting (FEC); and iv) determine the association between the presence of anemia and infection by these agents in a sheep herd from Bandeirantes City, Paraná State, southern Brazil.

5.3 Material and Methods

5.3.1 Ethical approval

This study was approved by the Ethics Committee for Animal Experimentation and Animal Welfare at Universidade Federal do Paraná (protocol n° 030/2019) and conducted according to the ethical principles of animal experimentation, adopted by the Brazilian College of Animal Experimentation.

5.3.2 Study

A total of 42 sheep from Bandeirantes County (23°06'28"S 50°21'36"W), Paraná State, southern Brazil, were evaluated for the presence of hemoplasmas, *Anaplasma* spp., piroplasms and GI parasites infection. Animals were co-grazed with cattle in a paddock, where tick infestation is common during all year.

5.3.3 Sampling

Sheep blood samples (up to 5 mL) were collected by venipuncture of the jugular vein using commercial tubes containing EDTA (BD Vacutainer®, Franklin Lakes, NJ, USA). Fecal samples were obtained directly by rectal collection, identified and stored in isothermal recipients until analysis.

Ticks found on animals were directly removed using a commercial hook (O'TOM Tick Twister®, Lavancia, France), and kept in absolute ethanol-labeled tubes until identification according to morphological taxonomic keys (Aragão and Fonseca, 1961; Guimarães et al., 2001).

5.3.4 Packed cell volume evaluation

The packed cell volume (PCV) was measured by routine centrifugation. A PCV value of <0.27 L/L was used as an indicator of anemia (Weiss et al., 2010). Thereafter, blood aliquots were stored at -20 °C until molecular testing.

5.3.5 Parasitological analysis

Fecal samples were analyzed by the eggs per gram (EPG) technique (Gordon & Whitlock 1939). Animals presenting EPG > 500 were considered positive (Ueno and Gonçalves, 1998).

5.3.6 DNA extraction

Isolation of genomic DNA from sheep blood samples was performed using a commercial kit (Illustra™ blood genomicPrep Mini Spin Kit, GE Healthcare, Little Chalfont, UK). Nuclease-free water was used in parallel as negative control to monitor cross-contamination.

5.3.7. Polymerase chain reactions (PCR)

DNA samples were submitted to the gene glyceraldehyde-3-phosphate dehydrogenase (GAPDH) housekeeping gene PCR, as previously described (Birkenheuer et al., 2003). GAPDH-positive samples were screened for *Mycoplasma* spp. by a PCR targeting a fragment

(900 bp) of the 16S rRNA gene of this bacterium (Hoelzle et al., 2011; Machado et al., 2017). Additionally, samples were also tested by PCR assays targeting a fragment (870 bp) of *msp4* gene of *A. ovis/A. marginale* (De La Fuente et al., 2007), and a fragment (551 bp) of the 18S rDNA gene of piroplasms (Almeida et al., 2012)

5.3.8 Sequencing

Amplicons obtained from four Mycoplasma-positive samples were purified by enzymatic purification (ExoSAP-IT™ PCR Product Cleanup Reagent, Thermo Scientific, Waltham, USA), evaluated by spectrophotometry for concentration and purity (NanoDrop™ One Spectrophotometer, Thermo Scientific, Waltham, USA), and sequenced in both directions by the Sanger method (3500 Genetic Analyzer, Applied Biosystems, Foster City, CA, USA). Thereafter, sequences were subjected to BLASTn analysis (Altschul et al., 1990) for determining the identity with the sequences deposited in the GenBank database. The nucleotide sequences of the *M. ovis* amplified herein were submitted to the GenBank® database (Genbank® accession numbers. XXX, XXX and XXX).

5.3.9 Phylogenetic analysis

Sequences were aligned using MAFFT 7.110 (Katoh and Standley, 2013). Best-fit evolutionary model was estimated as F81+I+G using jModeltest 2.1.4 (Darriba et al. 2012). The Bayesian Information Criterion (BIC) algorithm was used for phylogenetic inference. Three reconstruction was made with Fig.2 Tree 1.4.2. Polymorphisms were investigated in isolated sequences using DnaSP software v. 6.12.03.

5.4 Results

The mean PCV concentration for sheep was 0.30 L/L. Nine out of 42 (21.42%; 95% CI: 11.71-35.94%) animals were anemic. A total of 38 sheep were evaluated by EPG. Four out of 42 animals did not present feces at the time of sampling and were not evaluated. Twenty-four (63.15%; 95% CI: 47.2 – 76.6%) animals presented EPG values >500 and were considered positive for Strongylida type-eggs. A total of 24 adult tick specimens were collected from two/42 (4.76%; 95% CI: 1.32-1.57%) animals and all were identified as *R. microplus* species.

All samples consistently amplified the sheep GAPDH gene. Seven out of 42 (16.6%; 95% CI: 8.32–30.6%) animals were PCR-positive for *Mycoplasma* spp. Only one *Mycoplasma* spp. PCR-positive animal was anemic (PCV = 0.17L/L). The PCR results, parasitological analysis and presence of ticks has been summarized in Table 2. All samples tested negative to *Anaplasma* spp. and piroplasm species by PCR.

All *Mycoplasma*-positive samples sequenced showed $\geq 99\%$ identity with multiple *M. ovis* 16S rDNA gene sequences deposited in GenBank (accession nos. KU98374, KU983745, KU512718). The phylogeny showed two great clades dividing *Mycoplasma haemofelis* and *Mycoplasma suis* groups. Our sequences clustered together in a clade including *M. ovis* and *Mycoplasma* sp. sequences, inside the *M. suis* group. Polymorphism analysis revealed three different haplotypes from isolated sequences.

5.5 Discussion

Anemia is considered a concerning condition for animal health and production. Clinical manifestations may be represented by weakness, pallor of mucous membranes, hemoglobinuria and icterus. Regarding laboratorial findings, values of packed cell volume, total erythrocyte count and hemoglobin concentrations should be reduced (WEISS et al., 2010). In sheep, causatives of anemia are commonly linked to parasitological causes.

Herein we report the occurrence of *M. ovis* in sheep from Paraná State. Although there are no studies involving *M. ovis* occurrence in sheep from Brazil, our prevalence of 16.6% of infected animals is low when compared with data regarding *M. ovis* infection in different host species from Paraná State and Brazil. A study reporting *Mycoplasma* sp. occurrence in goats from northeastern region obtained prevalence of 39.30% (Machado et al., 2017). Regarding to wild animals, an overall prevalence of 58% of *M. ovis* infected deer from Brazilian southeast and midwestern regions is reported (Grazziotin et al., 2011a). In Paraná State, captive cervids have reached a prevalence of 87% of infection by *M. ovis* in the municipality of Foz do Iguaçu (Grazziotin et al., 2011b).

Infection by *M. ovis* in sheep may produce mild to severe anemia (Neimark et al., 2004). Our findings show that infection by *Mycoplasma* sp. in ovine may be often conducted as subclinical conditions, once only one positive animal presented anemia. Recently, a study conducted in Turkey showed that prevalence of hemotropic mycoplasma infection is lower in sheep with clinical signs than that classified as “healthy” (Aktas and Ozubek, 2017).

In the present work, we also evaluated the occurrence of GI parasites in sampled sheep. Concurrent infections with internal parasites may increase the pathogenicity of *M. ovis* in sheep (Ohtake et al., 2011). Infections by GI parasites and *M. ovis* in goats may lead animals to present anemia, pale mucous membrane and pasty feces (Abdullah et al., 2013). The only animal from our study that presented positive results for hemoplasmas and anemia obtained a low value of PCV (0.17 L/L). This animal has also obtained high values of EPG (1.700) from eggs identified as Strongylida type. The Strongylida suborder is known to shelter important parasites for ovine production, such as *Haemonchus contortus*, that may lead animals to severe anemic conditions (Amarante et al., 2014). Probably, the co-infection of GI parasites and hemoplasmas have worsened the animal health condition, which may represent an alert about hemoplasmas diagnosis in ovine herds with nematode infections occurrence.

The mean value of PCV obtained by *Mycoplasma* sp. infected sheep from the present study was 0.28 L/L, being within the reference values for this animal species (Weiss et al., 2010). Similar results were obtained in a study conducted in Argentina, where infected animals have mean value of PCV considered inside the normal range (Aguirre et al., 2009). Although sheep from our study were also co-infected with GI parasites, and clinical evaluation was unfortunately not performed, these results may suggest that these animals were well supporting the infections in relation to anemia development. Considering that eight hemoplasma negative animals obtained low PCV results, these results may be linked to other causes as other internal parasites or even hemoparasites do not target in this study.

Anaplasma marginale is a common tick-borne pathogen that infects primarily cattle from tropical and subtropical areas (Kocan et al., 2014). Recently, this bacterium was detected infecting sheep from Iran (Yousefi et al., 2017). Although there are no reports about *A. marginale* infection in sheep from Brazil, goats kept in a co-grazing system with cattle were found infected in the Northeastern region of the country (Da Silva et al., 2018). Although the main species vector involved in *A. marginale* transmission - *R. microplus* ticks - was found infecting animals from our study, all animals sampled presented negative results for *Anaplasma* spp. These results may suggest that acute infections caused by *Anaplasma* agents did not occur in the studied herd in the sampling moment.

The phylogenetic analysis showed that our sequences formed a cluster together with *M. ovis* sequences from different hosts and geographical origins. Sequences 3 and 5 from the present study clustered together with *M. ovis* isolated from Japanese serows (*Capricornis crispus*) from Japan (AB571119), meanwhile sequence 7 clustered together with *M. ovis*

isolated in sheep from Turkey (MF377458). All *M. ovis* sequences and *M. ovis*-related sequences formed a great cluster together within the *Mycoplasma suis* group.

Although our phylogenetic tree mostly presented satisfactory probabilities values, values regarding the study sequences presented low percentages. Polymorphism analysis showed three different haplotypes from the three sequences isolated. The presence of polymorphisms may cause incongruities in post probabilities values, although closely positions were obtained. Meanwhile, our phylogenetic results showed marked differences from sequences isolated in a same herd. Genotype diversity of *M. ovis* are also reported in flocks from China (Wang et al., 2017) and Mexico (Martinez-Hernandez et al., 2019).

The 16S rDNA gene is widely used for phylogenetic analysis, however characteristics as intra-genomic heterogenicities are considered a limiting factor (Rejendhran and Gunesakaran et al., 2011). In 2014, genome sequencing of *M. ovis* strain Michigan was reported showing the presence of two copies of the 16S rDNA gene (Deshuillers et al., 2014). Primers used in the present study are not capable of specifically targeting one or another copy from the *M. ovis*-16S rDNA which may represent unreliable data on sequence analysis.

Regarding the tick collection, one *R. microplus*-infested animal was also considered positive for *Mycoplasma* sp. Tick species *Rhipicephalus bursa* was implicated as a vector of *M. ovis* in small ruminants (Neimark et al., 2004). However, studies did not find significant association between tick presence and infection by *Mycoplasma* sp. in hosts (Rjeibi et al., 2015; Machado et al., 2017). The main vector of *Mycoplasma* sp. in sheep from Paraná remains unknown.

5.6 Conclusion

Anemia causes in the studied herd may be probably linked to GI infections. Different haplotypes of *Mycoplasma ovis* occurs in sheep from the north region of Paraná State.

5.7 References

Abdullah FFJ, Adamu L, Jamal MHAB, Osman AY, Haron AW, Awang DN, Roslim N. Parasitic Gastro-Enteritis (PGE) Concurrent with Eperythrozoonosis in a Goat: Case Report. *IOSR J Agricult Vet Sci* 2013; 4(1): 63-66.

- Aguirre DH, Thompson C, Neumann RD, Salatin AO, Gaido AB, Echaide ST. Brote de micoplasmosis clínica por *Mycoplasma ovis* en ovinos de Salta, Argentina. Diagnóstico clínico, microbiológico y molecular. *Rev Argent Microbiol* 2009; 41: 212-214.
- Alessandra T, Santo C. Tick-borne diseases in sheep and goats: Clinical and diagnostic aspects. *Small Rum Res* 2012; 106S: S6-S11.
- Aktas M, Ozubek S. A molecular survey of small ruminant hemotropic mycoplasmas in Turkey, including first laboratory confirmed clinical cases caused by *Mycoplasma ovis*. *Vet Microbiol* 2017; 208: 217-222.
- Almeida AP, Marcili A, Leite RC, Nieri-Bastos FA, Domingues LN, Martins JR, Labruna MB. *Coxiella* symbiont in the tick *Ornithodoros rostratus* (Acari: Argasidae). *Ticks Tick Borne Dis* 2012; 3(4): 203-6.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1999; 215: 403–410.
- Amarante AFT, Ragozo AMA, Silva BF. Os parasitos de ovinos. Editora Unesp Digital; 2014.
- Amarante AFT, Sales RO. Controle de Endoparasitoses de Ovinos: Uma Revisão. *Rev Bras Higiene San Anim* 2007; 1(2): 14-36.
- Birkenheuer AJ, Levy MG, Breitschewerdt EB. Development and evaluation of a seminested PCR for detection and differentiation of *Babesia gibsoni* (Asian genotype) and *B. canis* DNA in canine blood samples. *J Clin Microbiol* 2003; 41: 4172-4177.
- Aragão H, Fonseca F. Notas de ixolodologia. *Mem Inst Oswaldo Cruz* 1961; 59: 115-129.
- Boes KM, Goncarovs KO, Thompson CA, Halik LA, Santos AP, Guimaraes AMS, Feutz MM, Holman PJ, Vemulapalli R, Messick JB. Identification of a *Mycoplasma ovis*-like organism in a herd of farmed white-tailed deer (*Odocoileus virginianus*) in rural Indiana. *Vet Clin Pathol* 2012; 41(1): 77-83.
- Brito DRB, Santos ACG, Guerra RMSNC. Ectoparasitos em rebanhos de caprinos e ovinos na micro-região do Alto Mearim e Grajau, Estado do Maranhão. *Rev Bras Parasitol Vet* 2005; 14: 59– 63.
- Da Silva NB, Taus NS, Johnson WC, Mira A, Schnittger L, Valente JDM, Vidotto O, Masterson HE, Vieira TSWJ, Ueti MW, Vieira RFC. First report of *Anaplasma marginale* infection in goats, Brazil. *Plos One* 2018; 13(8): 30202140.
- Daddow KN. *Eperythrozoon ovis* – a cause of anemia, reduced production and decreased exercise tolerance in sheep. *Aust Vet J* 1979; 55: 605-606.

- De La Fuente J, Atkinson MW, Naranjo V, Mera IGF, Mangold AJ, Keating KA, Kocan KM. Sequence analysis of the *msp4* gene of *Anaplasma ovis* strains. *Vet Microbiol* 2007; 119(2-4): 375-381.
- Deshuillers PL, Santos AP, Nascimento NC, Hampel JA, Bergin IL, Dyson MC, Messick JB. Complete genome sequence of *Mycoplasma ovis* strain Michigan, a hemoplasma of sheep with two distinct 16S rRNA genes. *Genome Announc* 2014; 2(1): 1-2.
- Grazziotin AL, Duarte JMB, Szabó MPJ, Santos AP, Guimaraes AMS, Mohamed A, Vieira RFC, Barros-Filho IR, Biondo AW, Messick JB. Prevalence and Molecular Characterization of *Mycoplasma ovis* in Selected Free-Ranging Brazilian Deer Populations. *J Wild Dis* 2011a; 47(4): 1005-1011a.
- Grazziotin AL, Santos AP, Guimaraes AMS, Mohamed A, Cubas ZS, Oliveira MJ, Santos LC, Moraes W, Vieira RFC, Donatti L, Barros-Filho IR, Biondo AW, Messick JB. *Mycoplasma ovis* in captive cervids: Prevalence, molecular characterization and phylogeny. *Vet Microbiol* 2011b; 152: 415-419b.
- Guimarães JH, Battesti DMB, Tucci EC. Ectoparasitos de importância veterinária. Plêiade, 2001.
- Hampel JA, Spath SN, Bergin IL, Lim A, Bolin SR, Dyson MC. Prevalence and Diagnosis of Hemotropic *Mycoplasma* Infection in Research Sheep and Its Effects on Hematology Variables and Erythrocyte Membrane Fragility. *Comp Med* 2014; 64(6): 478-485.
- Hoelzle K, Winkler M, Kramer MM, Wittenbrink MM, Dieckmann SM, Hoelzle LE. Detection of *Candidatus Mycoplasma haemobos* in cattle with anemia. *Vet J* 2011; 187: 408-410.
- Hornok S, Hajtós I, Meli ML, Farkas I, Gönczi E, Meili T, Hofmann-Lehmann R. First molecular identification of *Mycoplasma ovis* and *Candidatus M. haemoovis* from goat, with lack of haemoplasma PCR-positivity in lice. *Acta Vet Hung* 2012; 60: 355–360.
- Hornok S, Meli ML, Erdos A, Hajtos I, Lutz H, Hoffman-Lehmann R. Molecular characterization of two different strains of haemotropic mycoplasmas from a sheep with fatal haemolytic anemia and concomitant *Anaplasma ovis* infection. *Vet Microbiol* 2009; 136: 372-377.
- IBGE. Pesquisa da Pecuária Municipal. Efetivos de rebanho, por tipo de rebanho, 2016. Available online in: <https://sidra.ibge.gov.br/tabela/3939> (Accessed in July 22, 2018).
- Johnson KA, Nascimento NC, Bauer AE, Weng H, Hammac GK, Messick JB. Detection of hemoplasma infection of goats by use of a quantitative polymerase chain reaction assay and risk factor analysis for infection. *Am J Vet Res* 2016; 77(8): 882-889.

- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 2013; 30(4): 772-780.
- Kocan KM, De La Fuente J, Blouin EF, Coetzee JF, Ewing SA. The natural history of *Anaplasma marginale*. *Vet Parasitol* 2010; 167: 95-107.
- Lane J, Jubb T, Shephard R, Webb-Ware J, Fordyce G. MLA Final Report: Priority List of Endemic Diseases for the Red Meat Industries. Meat and Livestock Australia, Sydney, Australia (2015). Available online in: <https://www.mla.com.au/research-and-development/search-rd-reports/final-report-details/Animal-Health-and-Biosecurity/Priority-list-of-endemic-diseases-for-the-red-meat-industries/2895>. (Accessed in March 13, 2019).
- Ma M, Chen Z, Liu A, Ren Q, Liu J, Liu Z, et al. Biological parameters of *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae) fed on rabbits, sheep, and cattle. *Korean J Parasitol*. 2016; 54: 301–305.
- Machado CAL, Vidotto O, Conrado FO, Santos NJR, Valente JDM, Barbosa IC, Trindade PWS, Garcia JL, Biondo AW, Vieira TSWJ, Vieira RFC. *Mycoplasma ovis* infection in goat farms from northeastern Brazil. *Comp Immunol Microbiol Infect Dis* 2017; 55: 1-5.
- R.G. Maggi, S.M. Compton, C.L. Trull, P.E. Mascarelli, B.R. Mozayeni, E.B. Breitschwerdt, Infection with hemotropic *Mycoplasma* species in patients with or without extensive arthropod or animal contact, *J. Clin. Microbiol.* 51 (2013) 3237–3241,
- Martinez-Hernández JM, Ballados-González GG, Fernández-Bandala D, Martínez-Soto S, Velázquez-Osorio V, Martínez-Rodríguez PB, Cruz-Romero A, Grostieta E, Lozano-Sardaneta Y, Colunga Salas P, Becker I, Sánchez-Montes S. Molecular detection of *Mycoplasma ovis* in an outbreak of hemolytic anemia in sheep from Veracruz, Mexico. *Trop Anim Health Product* 2019, 51(1): 243-248.
- Mason RW, Statham P. The determination of the level of *Eperythrozoon ovis* parasitaemia in chronically infected sheep and its significance to the spread of infection. *Animal Health Lab* 1991; 68(3): 115-116.
- Messick JB. Hemotrophic mycoplasmas (hemoplasmas): a review and new insights into pathogenic potential. *Vet Clinical Path* 2004; 33(1): 2-13.
- Neimark H, Hoff B, Ganter M. *Mycoplasma ovis* comb. nov. (formely *Eperythrozoon ovis*), an epierythrocytic agent of haemolytic anemia in sheep and goat. *Intern J Sys Evol Microbiol* 2004; 54: 365-371.
- Neitz WO, Alexander RA, Du Toit PJ. *Eperythrozoon ovis* (sp. nov.) Infection in Sheep. *Onderstepoort J Vet Sci Animal Indust* 1934; 3(2): 263-271.

- Ohtake Y, Nishizawa I, Sato M, Watanabe Y, Nishimura T, Matsubara K, Nagai K, Harasawa R. *Mycoplasma ovis* Detected in Free-Living Japanese Serows, *Capricornis crispus*. *J Vet Med Sci* 2011; 73(3): 371-373.
- Rajenhran J, Gunasekaran P. Microbial phylogeny and diversity: Small subunit ribosomal RNA sequence analysis and beyond. *Microbiol Res* 2011; 166: 99-110.
- Sheriff D, Clapp KH, Reid MA. *Eperythrozoon ovis* infection in south Australia. *Aust Vet J* 1966; 42: 169.
- Sykes JE, Lindsay LL, Maggi RG, Breitschwerdt EB. Human Coinfection with *Bartonella henselae* and Two Hemotropic *Mycoplasma* Variants Resembling *Mycoplasma ovis*. *J Clin Microbiol* 2010; 48(10): 3782-375.
- Tagawa M, Takeuchi T, Fujisawa T, Konno Y, Yamamoto S, Matsumoto K, Yokoyama N, Inokuma H. A Clinical Case of Severe Anemia in a Sheep Coinfected with *Mycoplasma ovis* and ‘*Candidatus Mycoplasma haemovis*’ in Hokkaido, Japan. *Intern Med* 2012; 74(1): 99-102.
- Taylor MA, Coop RL, Wall RL. *Veterinary Parasitology*, 3th ed., Wiley-Blackwell, Ames, Iowa, 2007.
- Ueno H, Gonçalves PC. *Manual para Diagnóstico das Helmintoses de Ruminantes*. 4 ed. Tóquio: Japan International Cooperation Agency, 1998.
- Wang X, Cui Y, Zhang Y, Shi K, Yan Y, Jian F, Zhang L, Wang R, Ning C. Molecular characterization of hemotropic mycoplasmas (*Mycoplasma ovis* and ‘*Candidatus Mycoplasma haemovis*’) in sheep and goats in China. *BMC Vet Res* 2017; 13: 142-150.
- Weiss DJ, Wardrop KJ, Schalm OW. *Schalm’s Veterinary Hematology*, 4th ed., Wiley-Blackwell, Ames, Iowa, 2010.
- Yeruham I, Hadani A, Galker F. Some epizootiological and clinical aspects of ovine babesiosis caused by *Babesia ovis* – a Review. *Vet Parasitol* 1998; 74: 153-163.
- Yousefi A, Rahbari S, Shayan P, Sadeghi-dehkordi Z, Bahonar A. Molecular detection of *Anaplasma marginale* and *Anaplasma ovis* in sheep and goat in west highland pasture of Iran. *Asian Pac J Trop Biomed* 2017; 7:455-459.



Figure 1. Phylogenetic tree based on partial sequences of the 16S rDNA gene, showing the relationship between the *Mycoplasma* sp. detected in sheep from this study and other hemoplasmas by Bayesian Inference. *Bacillus subtilis* and *Mycoplasma bovis* were used as outgroups. Analyses were carried out applying the F81+I+G model and 1000 bootstrap replicates for all analyses.

Table 2. Obtained results regarding PCV values, amplification of the *Mycoplasma* spp. 16S rRNA fragment by PCR, EPG values and tick presence/identification in sheep from the present study.

Animal identification	PCV value	Mycoplasma PCR result	Parasitological analysis	Presence of <i>R. microplus</i> ticks
1	29,9	+	7.950 EPG	
2	32,3	+	250 EPG	
3	23,9		Negative	
4	28,1		450 EPG	
5	27,0		300 EPG	
6	31,6	+	450 EPG	
7	27,4	+	Negative	1 male and 10 females
8	21,5		Not collected	
9	32,4		Negative	
10	29,1		1.750 EPG	
11	32,7		Negative	
12	28,6		Negative	
13	21,9		5.200 EPG	
14	31,6		900 EPG	
15	19,3		2.750 OPG	
16	30,5		50 EPG	
17	34,4		1.550 EPG	
18	23,6		3.400 EPG	
19	34,4		350 EPG	
20	34,3		Not collected	
21	33,7	+	2.000 EPG	
22	26,9		3.500 EPG	
23	29,1	+	Not collected	
24	29,2		400 EPG	
25	38,1		150 EPG	
26	20,9		800 EPG	1 male and 12 females
27	29,8		1.450 EPG	
28	31,5		950 EPG	
29	29,8		2.150 EPG	
30	31,6		500 EPG	
31	28,9		3.400 EPG	
32	35,8		1.200 EPG	
33	33,1		2.150 EPG	
34	15,0		Not collected	
35	34,0		Not collected	
36	33,5		500 EPG	
37	32,1		2.950 EPG	
38	27,3		4.500 EPG	
39	28,2		7.000 EPG	
40	17,6	+	4.550 EPG	
41	29,5		1.050 EPG	
42	36,1		1.700 EPG	

6 APPENDIX: SECOND MANUSCRIPT FOR SUBMISSION

A parallel research made during the master's course.

Title: Hemotrophic hemoplasma infection in Rhesus Monkeys (*Macaca mulatta*) from a research colony in Rio De Janeiro, Brazil.

6.1 Introduction

The Mollicutes class involves about of 200 classified species of bacteria. Among them, the ones comprised in the *Mycoplasma* genus are distinguished by cell wall lack (Razin et al., 1998; Trachtenberg, 2005). However, this characteristic did not prevent *Mycoplasma* to infect a wide range of host species including mammals, birds, arthropods and plants. With a notable small genotype, these species become desirable models for genomics studies (Dybvig and Voelker, 1996).

Since few decades ago, species belonging to *Haemobartonella* and *Eperythrozoon* genus, from Anaplasmataceae family, were proposed to be re-classified into the *Mycoplasma* genus in order to reflect its phylogenetic relationship obtained by studies with the 16S rRNA gene from these species (Rikihisa et al., 1997; Neimark et al., 2001).

Such studies allowed the classification of some *Mycoplasma* species as hemotrophic *Mycoplasmas*, also known as hemoplasmas. These species are considered small pleomorphic bacteria that parasitize the surface of red blood cells from different vertebrate hosts and may lead them to anemia conditions (Messick et al., 2004). Hemoplasma infection is widely studied in a large range of animal hosts, both domestic and wild species.

The use of animal experimentation in biomedical researches had vastly increased the scientific knowledge and human health (Institute of Medicine and Nacional Research Council, 1998). Health monitoring programs in laboratory kept animals are essential once only animals free of pathogens that may modify its physiological parameters are considered suitable for research uses (Baker, 1998).

Although mucous *Mycoplasma* infections are well documented in laboratory animals, few studies are related to hemoplasmas. In research sheep, hemoplasmas caused subclinical and acute infections, raising concerns about the suitability of these animals for biomedical studies (Hampel et al., 2014). *Mycoplasma coccoides* (formerly *Eperythrozoon coccoides*) is

considered a common species that may cause latent infection in laboratory rodents (Glasgow et al., 1974). This species is reported as a cause of influence in late studies involving other hemoparasites infections such as *Plasmodium chabaudi* (Ott et al., 1967), *Trypanosoma brucei* (Molyneux, 1970) and *Plasmodium berghi* (Andrade Jr et al., 1986). Moreover, *M. coccoides* showed to interfere in different study fields, as host interferon production against viral infections (Glasgow et al, 1971) and significance factor in tumor studies (Nelson, 1955).

Mycoplasma muris (formerly *Haemobartonella muris*) is also well documented as a causative of latent infection in laboratory mouse (Bartlett and Pease, 1974). This species may cause synergism mechanisms in concurrent infections with *P. berghi* causing activation of latent *M. muris* with intense infection rates (Hsu and Geiman, 1952).

Regarding non-human primates (NHP), it is estimated that 100,000 to 200,000 animals are annually used in research worldwide (Carlsson et al., 2004). These animals are described as hosts of three hemoplasmas species that remain partial characterized: ‘*Candidatus Mycoplasma kahanei*’ (Neimark et al., 2002), ‘*Candidatus Mycoplasma aoti*’ (Barker et al., 2011) and ‘*Candidatus Mycoplasma haemomacaque*’ (Maggi et al., 2013).

The present study aims to report *Mycoplasma* sp. infection detected in blood samples from rhesus monkey (*Macaca mulatta*) from a research center in Brazil.

6.2 Material and Methods

6.2.1 Sampling

In 2018, blood samples from rhesus monkeys kept in captivity for laboratory uses were collect twice by venipuncture and stored in tubes containing EDTA. The laboratory colony is maintained at a certified Research Center in Rio de Janeiro City, southeastern Brazil for use on studies focusing on arboviruses as Yellow Fever and Zika.

The first collection was performed in February and a total of eight animals were sampled. Second collection were performed in the same animals in December and only six animals were sampled once two of them died.

Animals did not present ectoparasites at the time of sampling. Researchers stated that all non-human primates receive a preventive application of ivermectin monthly.

6.2.2 DNA extraction and polymerase chain reaction (PCR) assays

DNA was obtained from 200 uL of blood from each sample using a commercial kit (Illustra™ blood GenomicPrep Mini Spin Kit, GE Healthcare, Little Chalfont, UK). A conventional PCR for the non-human primate house-keeping gene glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was performed in all samples to ensure successful DNA extraction (Birkenheuer et al., 2003). Samples were screened for hemoplasmas using previously described pan-hemoplasma primers targeting the 16S rDNA gene (~900 bp) (Hoelze et al., 2001; Machado et al., 2017). A cat-positive blood sample for '*Candidatus M. haemominutum*' and ultrapure water were used as positive and negative controls, respectively.

6.2.3 Sequencing and phylogenetic analysis

Amplicons obtained from three samples were purified by enzymatic purification (ExoSAP-IT™ PCR Product Cleanup Reagent, Thermo Scientific, Waltham, USA), evaluated by spectrophotometry for concentration and purity (NanoDrop™ One Spectrophotometer, Thermo Scientific, Waltham, USA), and sequenced in both directions by Sanger method (3500 Genetic Analyzer, Applied Biosystems, Foster City, CA, USA). Thereafter, sequences were subjected to BLASTn analysis (Altschul et al., 1990) for determining the identity with sequences deposited in GenBank database. Sequences were then aligned using MAFFT (Katoh et al., 2013), and improved on GUIDANCE2 (Sela et al., 2015). Best-fit evolutionary model was estimated as F81+I+G by jModeltest 2.1.4 (Darriba et al. 2012). The Bayesian Information Criterion (BIC) algorithm was used for phylogenetic inference and made by BEAST 1.8 with three independent runs of 50.000.000 MCMC. Trees were combined with TreeAnnotator 1.8 software and reconstruction was visualized with FigTree 1.4.0 software.

6.3 Results

All samples from both samplings successfully amplified the GAPDH gene and were submitted to *Mycoplasma* sp. detection. From the first set of sample collection, a total of five/8 (62,5%; CI: 3.05 – 8.63) rhesus monkeys tested positive for *Mycoplasma* spp. One sample that presented a strong band on electrophoresis was chose for sequencing and generated a 715 bp

sequence that was deposited in GenBank (XXXX.X). When analyzed by BLASTn the sequence showed similarity of 99% with ‘*Ca. M. haemomacae*’ sequences isolated from cynomolgus monkeys (*Macaca fascicularis*) in the United States (KC512401.1; AB820288.1).

Regarding the second sampling, three/6 (50%; CI: 1.87 – 8.12) animals tested positive for *Mycoplasma* spp. From these animals, one was negative in the first sampling. Two samples presenting strong bands on electrophoresis were chosen for sequencing and generated sequences of 1.326 bp and 976 bp that were deposited in GenBank with accession number XXXX.X and XXXX.X, respectively.

In phylogeny analysis, our sequence formed a clade with ‘*Ca. M. haemomacae*’ isolates from USA (KC512401) and Japan (AB820288) within the *M. haemofelis* group (EU442639).

6.4 Discussion

Infection by *Mycoplasma* sp. was already described in rhesus monkeys (Peters et al., 1974). Based on partial 16S rRNA amplification and phylogeny, our results suggest that the species involved in the described infection is ‘*Ca. M. haemomacae*’. This hemoplasma was recently proposed after its isolation in research cynomolgus monkeys (*Macaca fascicularis*) from USA through analysis of the 16S rDNA and partial RNase P genes (Maggi et al., 2013). Few later, this species was reported infecting captive *Macaca fuscata* from Japan with closely phylogenetic relationship with ‘*Ca. M. haemomacae*’ isolated in the USA (Maggi et al., 2013; Sashida et al., 2013).

A closely genotype was reported infecting free-range and captive monkeys from *Sapajus* genus in two different regions of northeastern Brazil and was suggested as a possible novel species of hemoplasma (Bonato et al., 2015; Ramalho et al., 2017). Otherwise, our study describes a genotype very close to ‘*Ca. M. haemomacae*’ from USA and Japan (Maggi et al., 2013; Sashida et al., 2013) infecting a research colony of monkeys in southeast Brazil.

Prevalence from the present study (62,5%) is lower when compared to studies that investigate hemoplasma infection in monkeys from research colonies out of Brazil, which ranged from 84.6% to 100% (Dillberger et al., 1974; Maggi et al., 2013; Sashida et al., 2014). However, it was high compared with studies conducted in Brazil with captive and free-range NHP that presented 35.7% and 25% rates (Bonato et al., 2015; Cubilla et al., 2017). Although it is more expected free-range animals to be infected, hemoplasma infection rates may be also high in colonies of laboratory animals (Conrado et al., 2015).

Regarding phylogenetic analysis, sequence 5 formed a robust clade together with ‘*Ca. M. haemomacaque*’ isolates from USA and Japan (Maggi et al., 2013; Sashida et al., 2013), suggesting a greater proximity to this species. On the other hand, our sequence clustered distantly from *Mycoplasma* sp. genotypes isolated in *Sapajus* spp. monkeys from Brazil. Thus, a hypothesis that a potentially novel hemoplasma species circulates in NHP from northeastern region of Brazil, differently from the one isolated herein, may be raised. However, larger gene fragments and other genomic regions should be analyzed to confirm this statement.

Sequences from animals 2 and 4 formed a separately clade from animal 5 sequence and ‘*Ca. M. haemomacaque*’ isolates from USA and Japan (Maggi et al., 2013; Sashida et al., 2013) clade, although it greatly presents posterior probabilities values. This finding may suggest a homology from the sequences isolated in the present study and differences may be due to sequencing procedures.

Mycoplasma genotype close to ‘*Ca. M. kahanei*’ is reported infecting *Alouatta* sp. monkeys from southern Brazil (Cubilla et al., 2017) and may be circulating in *Sapajus* monkeys from Brazilian Amazon (Bonato et al., 2015). Our phylogenetic analysis confirmed this relationship, however it clustered in the same group with *M. suis* (EU603330) whereas our *Mycoplasma* genotype was within the *M. haemofelis* group, suggesting that at least three genetically distinct hemoplasmas occur in NHP from Brazil.

Studies based on entire 16S rRNA gene and the intergenic spacer region between 16S and 23S rDNA genes showed that a monkey colony was entirely infected by the same strain of ‘*Ca. M. haemomacaque*’ in Japan (Sashida et al., 2013). Although we obtained a high prevalence in our study (5/8; 62.5%), unfortunately it was not possible to sequence more than one positive sample to identify diversity between the isolates.

Screening of research animals by hemoplasma-PCR assays may be a desirable tool to avoid complications caused by infections (Hampel et al., 2014). Hematological or biochemical abnormalities were not associated to ‘*Ca. M. haemomacaque*’ or other related genotypes in *Macaca fascicularis* and *Sapajus* sp. monkeys (Maggi et al., 2013; Ramalho et al., 2017). However, a moderate drop of erythrocytes may occur in splenectomized *Saimiri sciureus* monkeys (Contaim and Michel, 1999) and anemic episodes leading to death were reported in *M. fascicularis* infected by *Mycoplasma* sp. (Dillberger et al., 1994).

Mycoplasma infection may lead to hematological changes (Hampel et al., 2014) and haptoglobin raising in apparently healthy hosts (Vilhena et al., 2018). Thus, outwardly “normal” animals may be not appropriate for researches because of unapparent infection effects.

Moreover, when infections are clinically visible, the use of drugs to clear the pathogens may influence on animal parameters and research results (Baker et al., 1999).

A late study regarding *M. coccoides* influence on viral infections showed that this hemoparasite species may influence rediculoendothelial system causing suppression of interferon response in rats (Glasgow et al., 1971). Once the animals sampled in the present study are used for arboviruses studies, it is important to investigate if hemoplasmas contamination may represent a bias in research development.

Transmission of *Mycoplasma* sp. agents in research groups of monkeys may be associated with needles or contaminated nasogastric tubes (Dillberger et al., 1994). Some *Mycoplasma* infections are associated with vectors transmissions, as ticks and fleas (Neirmark et al., 2004; Woods et al., 2005). Once the animals sampled herein received a monthly application of ivermectin, transmission may be associated with experimental manipulation or direct contact between animals. Also, studies involving *M. muris* showed that latent infections may occur in laboratory animals with absence of ectoparasites (Bartelett and Pease, 1974). Transplacental transmission possibility is also concerned (Neimark et al., 2002). In all cases, it reveals the importance of ensure biosecurity proceedings and preventive screenings aiming to avoid contamination spread in research monkey colonies.

Moreover, some *Mycoplasma* species were described zoonotically infecting humans (Santos et al., 2008; Bosnoc et al., 2009; Sykes et al., 2010). Zoonotic potential should always be concerned to highlight possibilities of an occupational infection, especially for immunosuppressed persons.

6.5 Conclusion

Monkeys from the studied research colony are infected by *Mycoplasma* sp. More studies are necessary to elucidate the occurrence of hematological or clinical changes in infected animals and its transmission.

6.6 References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. *J Mol Biol* 1999; 215: 403–410.

- Andrade Jr HF, Laurenti MD, Goto H, Duarte MIS, Corbett P. Malária experimental: contaminação de cepas e animais de biotério por *Eperythrozoon coccoides*. *Rev Inst Med Trop São Paulo* 1986; 28(2): 246-252.
- Baker DG. Natural Pathogens of Laboratory Mice, Rats and Rabbits and Their Efforts on Research. *Clin Microbiol Rev* 1998; 11(2): 231-266.
- Barker EN, Helps CR, Neimark H, Peters IR, Peters W, Tasker S. A novel haemoplasma species identified in archived primate blood smears. *Vet Microbiol* 2011; 149: 478-481.
- Birkenheuer AJ, Levy MG, Breitschewerdt EB. Development and evaluation of a seminested PCR for detection and differentiation of *Babesia gibsoni* (Asian genotype) and *B. canis* DNA in canine blood samples. *J Clin Microbiol* 2003; 41: 4172-4177.
- Bonato L, Figueiredo MAP, Gonçalves LR, Machado RZ, André MR. Occurrence and molecular characterization of Bartonella sp. and hemoplasmas in neotropical primates from Brazilian Amazon. *Comp Immunol Microbiol Infect Dis* 2015; 42: 15-20.
- Bosnoc D, Baresic M, Branimir A, Sentic M, Cerovec M, Mayer M, Cikes N. Rare zoonosis (hemotropic mycoplasma infection) in a newly diagnosed systemic lupus erythematosus patient followed by a Nocardia asteroides pneumonia. *Braz J Infect Dis* 2010; 14(1): 92-95.
- Carlsson HE, Schapiro SJ, Farah I, Hau J. Use of Primates in Research: A Global Overview. *American J Primatol* 2004; 63: 225-237.
- Ceola CF, Sampaio J, Blatt SL, Cordova CMM. *Mycoplasma* infection and inflammatory effects on laboratory rats bred for experimental research. *Rev Pan-Amaz Saude* 2016; 7(2): 59-66.
- Conrado FO, Nascimento NC, Santos AP, Zimpel CK, Messick JB, Biondo AW. Occurrence and identification of hemotropic mycoplasmas (Hemoplasmas) in free ranging and laboratory rats (*Rattus norvegicus*) from two Brazilian zoos. *BMC Vet Res* 2015; 11: 286-296.
- Contamin H, Michel JC. Haemobartonellosis in Squirrel Monkeys (*Saimiri sciureus*): Antagonism between *Haemobartonella* sp. and Experimental Plasmodium falciparum Malaria. *Experimen Parasitol* 1999; 91: 197-305.
- Cubilla MP, Santos LC, Moraes W, Cubas ZS, Leutenegger M, Estrada M, Vieira RFC, Soares MJ, Lindsay LL, Sykes JE, Biondo AW. Occurrence of hemotropic mycoplasmas in non-human primates (*Alouatta caraya*, *Sapajus nigritus* and *Callithrix jacchus*) of southern Brazil. *Comp Immunol Microbiol Infec Dis* 2017; 52: 6 – 13.
- Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. *Nat Methods* 2012; 9: 772.

- Dillberger JE, Loudy DE, Adler RR, Gass JH. Hemobartonella-like Parasites in Cynomolgus Monkeys (*Macaca fascicularis*). *Vet Pathol* 1994; 31(3): 301-307.
- Dybvig K, Voelker LL. Molecular Biology of mycoplasmas. *Ann Rev Microbiol* 1996; 50: 25-27.
- Glasgow LA, Murrer AT, Lombardi PS. *Eperythrozoon coccoides*. II. Effect on Interferon Production and Role of Humoral Antibody in Host Resistance. *Infection and Immunity* 1974; 9(2): 266-272.
- Glasgow LA, Odugbemi T, Dwyer P, Ritterson AL. *Eperythrozoon coccoides*. I. Effect on the Interferon Response in Mice. *Infection and Immunity* 1971; 4(4): 425-430.
- Hampel JA, Spath SN, Bergin IL, Lim A, Bolin SR, Dyson MC. Prevalence and Diagnosis of Hemotrophic *Mycoplasma* Infection in Research Sheep and Its Effects on Hematology Variables and Erythrocyte Membrane Fragility. *Comparative Med* 2014; 64(6): 478-485.
- Hoelzle K, Winkler M, Kramer MM, Wittenbrink MM, Dieckmann SM, Hoelzle LE. Detection of *Candidatus Mycoplasma haemobos* in cattle with anemia. *Vet J* 2011; 187: 408-410.
- Institute of Medicine and National Research Council. 1998. Use of Laboratory Animals in Biomedical and Behavioral Research. Washington, DC: The National Academies Press.
- Hsu DYM, Geiman QM. Synergistic effect of *Haemobartonella muris* on *Plasmodium berghei* in white rats. *American J Trop Med Hyg* 1952; 1(5): 747-760.
- Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* 2013; 30(4): 772-780.
- Machado CAL, Vidotto O, Conrado FO, Santos NJR, Valente JDM, Barbosa IC, Trindade PWS, Garcia JL, Biondo AW, Vieira TSWJ, Vieira RFC. *Mycoplasma ovis* infection in goat farms from northeastern Brazil. *Comp Immunol Microbiol Infect Dis* 2017; 55: 1-5.
- Maggi RG, Mascarelli PE, Balakrishnan N, Rohde CM, Kelly CM, Ramaiah L, Leach MW, Breitschwerdt EB. “*Candidatus Mycoplasma haemomacaque*” and *Bartonella quintana* bacteremia in cynomolgus monkeys. *J Clin Microbiol* 2013; 51(5): 1408-1411.
- Messick JB. Hemotrophic mycoplasmas (hemoplasmas): a review and new insights into pathogenic potential. *Vet Clinical Path* 2004; 33(1): 2-13.
- Molyneux DH. Relationship between *Eperythrozoon coccoides* and *Trypanosoma* (Trypanozoon) *brucei brucei* in experimentally infected mice. *Annals Trop Med Parasitol* 1970; 64(3): 325-328.

- Neimark H, Barnaud A, Gounon P, Michel JC, Contamin H. The putative *haemobartonella* that influences *Plasmodium falciparum* parasitaemia in squirrel monkeys is a haemotrophic mycoplasma. *Microbes Infect* 2002; 4: 693-698.
- Neimark H, Hoff B, Ganter M. *Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an eperythrocyclic agent of haemolytic anemia in sheep and goat. *Intern J Sys Evol Microbiol* 2004; 54: 365-371.
- Neimark H, Johansson KE, Rikihisa Y, Tully JG. Proposal to transfer some members of the genera *Haemobartonella* and *Eperythrozoon* to the genus *Mycoplasma* with descriptions of 'Candidatus *Mycoplasma haemofelis*', 'Candidatus *ycoplasma haemomuris*', 'Candidatus *Mycoplasma haemosuis*' and 'Candidatus *Mycoplasma wenyonii*'. *Inter J Sys Evolution Microbiol* 2001; 51: 891-899.
- Nelson, JB. An ascites tumor appearing during the passage of *Eperythrozoon coccoides* in mice. *J Exp Med* 1956; 103(6): 743-752.
- Ott KJ, Astin JK, Leslie AS. *Eperythrozoon coccoides* and Rodent Malaria: *Plasmodium chabaudi* and *Plasmodium berghei*. *Experimental Parasitology* 1967; 21: 68-77
- Peters W, Molyneux DH, Howells RE: *Eperythrozoon* and *Haemobartonella* in monkeys. *Ann Trop Med Parasitol* 1974; 68: 47-50.
- Ramvalho AC, Guerra RR, Mongrue ACB, Vidotto O, Lucena RB, Guerra MVSF, Vieira TSWJ, Vieira RFC. *Mycoplasma* sp. infection in captive Marcgrave's capuchin monkeys (*Sapajus flavius*). *Comp Immunol Microbiol Infect Dis* 2017; 51: 34-36.
- Razin S, Yogev D, Naot Y. Molecular Biology and Pathogenicity of Mycoplasmas. *Microbiol Mol Biol Rev* 1998; 62(4): 1094-1156.
- Rikihisa Y, Kawahara M, Wen B, Kociba G, Fuerst P, Kawamori F, Suto C, Shibata S, Futohashi M. Western Immunoblot Analysis of *Haemobartonella muris* and Comparison of the 16S rRNA Gene Sequences of *H. muris*, *H. felis* and *Eperythrozoon suis*. *J Clin Microbiol* 1997; 35(4): 823-829.
- Santos AP, Santos RP, Biondo AW, Dora JM, Goldani LZ, Oliveira ST, Guimarães AMS, Timenetsky J, Morais HA, González FHD, Messick JB. Hemoplasma infection in HIV-positive Patient, Brazil. *Emerg Infect Dis* 2008; 14(12): 1922-1924.
- Sashida H, Suzuki Y, Rokuhara S, Nagai K, Harasawa R. Molecular Demonstration of Hemotropic Mycoplasmas in Wild Japanese Monkeys (*Macaca fuscata*). *J Vet Med Sci* 2014; 76(1): 97-101.

Sela I, Ashkenazy H, Katoh K, Pupko T. GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. *Nuclei Acid Res* 2015; 43: W7-W14.

Sykes JE, Lindsay LE, Maggi RG, Breitschwerdt E. Human Coinfection with *Bartonella henselae* and Two Hemotropic Mycoplasmas Variants Resembling *Mycoplasma ovis*. *J Clin Microbiol* 2010; 48(10): 3782-3785.

Trachtenberg S. Quick Guide: Mollicutes. *Current Biol* 2005; 15(13): PR483-R484.

Vilhena H, Tvarijonaviciute A, Céron JJ, Ramiro Pastorinho M, Martinez-Subiela S, Pastor J, Silvestre-Ferreira AC. Acute phase proteins response in cats naturally infected by hemotropic mycoplasmas. *Comp Immunol Microbiol Infect Dis* 2018; 56: 1-5.

Woods JE, Brewer MM, Hawley JR, Misnewski N, Lappin MR. Evaluation of experimental transmission of *Candidatus Mycoplasma haemominutum* and *Mycoplasma haemofelis* by *Ctenocephalides felis* to cats. *Am J Vet Res* 2005; 66(6): 1008-1012.




Figure 2. Phylogenetic tree based on partial 16S rDNA gene fragments of *Mycoplasma* sp. Tree was constructed by Bayesian Inference and a sequence from *Bacillus subtilis* was used as outgroup. Analysis were carried out applying the F81+I+G model and 1000 bootstrap replicates for all analyses. Our sequences are represented by ** after its name.

7 FINAL CONCLUSIONS

The clinical findings linked to anemia that were affecting the studied herd may be linked to infection by GI parasites. Sheep from the present study were parasitized by *R. microplus* ticks. The mean PCV concentration for sheep was 0.30 L/L. Nine out of 42 (21.42%; 95% CI: 11.71-35.94%) animals were anemic. Seven out of 42 (16.6%; 95% CI: 8.32–30.6%) animals were PCR-positive for *Mycoplasma* spp. None of the animals were positive for *Anaplasma* spp. and piroplasmids.

Twenty-four (63.15%; 95% CI: 47.2 – 76.6%) animals presented EPG values >500 and were considered positive for Strongylida type-eggs. All *Mycoplasma*-positive samples sequenced showed $\geq 99\%$ identity with multiple *M. ovis* 16S rDNA gene sequences deposited in GenBank (accession nos. KU98374, KU983745, KU512718). The phylogeny showed two great clades dividing *Mycoplasma haemofelis* and *Mycoplasma suis* groups. Our sequences clustered together in a clade including *M. ovis* and *Mycoplasma* sp. sequences, inside the *M. suis* group. Polymorphism analysis revealed three different haplotypes from isolated sequences.

REFERENCES

- ABDULLAH, J.; ADAMU, L.; JAMAL, M.H.A.B.; OSMAN, A.Y.; et al. Parasitic Gastro-Enteritis (PGE) Concurrent with Eperythrozoonosis in a Goat: A Case Report. **IOSR Journal of Agriculture and Veterinary Science**, v. 4, n. 1, p. 63-66, 2013.
- AGUIRRE, D.H.; THOMPSON, C.; NEUMANN, R.D.; SALATIN, A.O.; et al. Brote de micoplasmosis clínica por *Mycoplasma ovis* en ovinos de Salta, Argentina. Diagnóstico clínico, microbiológico y molecular. **Revista Argentina de Microbiología**, v. 41: 212-214, 2009.
- AKTAS, M.; OZUBEK, S. A molecular survey of small ruminant hemotropic mycoplasmas in Turkey, including first laboratory confirmed clinical cases caused by *Mycoplasma ovis*. **Veterinary Microbiology**, v. 208, p. 217-222, 2017.
- AKTAS, M.; OZUBEK, S. *Anaplasma ovis* genetic diversity detected by major surface protein 1a and its prevalence in small ruminants. **Veterinary Microbiology**, v. 217, p. 13–17, 2018.
- AKTAS, M.; OZUBEK, S. Short communication A molecular survey of small ruminant hemotropic mycoplasmosis in Turkey, including first laboratory confirmed clinical cases caused by *Mycoplasma ovis*. **Veterinary Microbiology**, v. 208, p. 217–222, 2017.
- ALANI, A. J.; HERBERT, I. V. Pathogenesis of Infection with *Theileria recondita* (Wales) Isolated from *Haemaphysalis punctata* from North Wales. **Veterinary Parasitology**, v. 28, p. 293–301, 1988.
- ALAZANI, A.D.; SAID, E.A.; GHONEIM, A.M.; et al. A comprehensive evaluation and first molecular report of *Theileria ovis* infection in small ruminants in Saudi Arabia. **Tropical Animal Health and Production**, v. 51, n. 1, p. 89-98, 2018.
- ALESSANDRA, T.; SANTO, C. Tick-borne diseases in sheep and goats: Clinical and diagnostic aspects. **Small Ruminant Research**, v. 106S, p. S6-S11, 2012.
- ALI, A. M.; SALIH, D. A.; NJAHIRA, M. N.; et al. Genotyping of *Theileria lestoquardi* from sheep and goat in Sudan to support. **Veterinary Parasitology**, v. 239, p. 7–14, 2017.
- ALLSOPP, B. A. Natural history of *Ehrlichia ruminantium*. **Veterinary Parasitology**, v. 167, p. 123–135, 2010.
- ALMEIDA, A.P.; MARCILI, A.; LEITE, R.C.; NIERI-BASTOS, F.A.; DOMINGUES, L.N.; et al. *Coxiella* symbiont in the tick *Ornithodoros rostratus* (Acari: Argasidae). **Ticks and Tick-Borne Diseases**, p. 3, v.4, p. 203-6, 2012.
- ALTAY K., DUMANLI, N.; AKTAS, M. A study on ovine tick-borne hemoprotozoan parasites (*Theileria* and *Babesia*) in the East Black Sea Region of Turkey. **Parasitology Research**, v. 111, p. 149-153, 2012.

- ALTAY, K.; DUMANLI, N.; HOLMAN, P. J.; AKTAS, M. Detection of *Theileria ovis* in naturally infected sheep by nested PCR. **Veterinary Parasitology**, v. 127, p. 99–104, 2005.
- ALTSCHUL, S.F.; GISH, W.; MILLER, W.; MYERS, E.W.; LIPMAN, D.J. Basic local alignment search tool. **Journal of Molecular Biology**, v. 215, p. 403–410, 1999.
- AMARANTE, A.F.T.; RAGOZO, A.M.A.; SILVA, B.F. **Os parasitos de ovinos**. 1a Edição. São Paulo: Editora Unesp Digital; 2014.
- AMARANTE, A.F.T.; SALES, R.O. Controle de Endoparasitoses de Ovinos: Uma Revisão. **Revista Brasileira de Higiene e Sanidade Animal**, v. 1, n. 2, p. 14-36, 2007.
- ANDRADE JR, H.F.; LAURENTI, M.D.; GOTO, H.; DUARTE, M.I.S.; CORBETT, P. Malária experimental: contaminação de cepas e animais de biotério por *Eperythrozoon coccoides*. **Revista do Instituto de Medicina Tropical de São Paulo**, p.28, v.2, p. 246-252, 1986.
- AOUADI, A.; LEULMI, H.; BOUCHEIKHCHOUKH, M.; BENAKHLA, A. Molecular evidence of tick-borne hemoprotozoan-parasites (*Theileria ovis* and *Babesia ovis*) and bacteria in ticks and blood from small ruminants in Northern Algeria. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 50, p. 34–39, 2017.
- ARAGÃO, H.; FONSECA, F. Notas de ixolodologia. **Memórias do Instituto Oswaldo Cruz**, v. 59, p. 115-129, 1961.
- AYDIN, M.F.; AKTAS, M.; DUMANLI, N. Molecular Identification of *Theileria* and *Babesia* in sheep and goats in the Black Sea Region in Turkey. **Parasitology Research**, v. 112, n. 8, p. 2817-2824, 2013.
- BAKER, D.G. Natural Pathogens of Laboratory Mice, Rats and Rabbits and Their Effects on Research. **Clinical Microbiology Review**, p. 11, v. 2, p. 231-266, 1998.
- BARKER, E.N.; HELPS, C.R.; NEIMARK, H.; PETERS, I.R.; PETERS, W.; et al. A novel haemoplasma species identified in archived primate blood smears. **Veterinary Microbiology**, v. 149, p. 478-481, 2011.
- BAZMANI, A.; ABNOLHOOSHYAR, A.; IMANI-BARAN, A.; et al. Semi-nested polymerase chain reaction-based detection of *Babesia* spp. in small ruminants from Northwest of Iran. **Veterinary World**, v. 11, n. 3, p. 268-273, 2018.
- BEKKER, C. P. J.; VOS, S. DE; TAOU, A.; SPARAGANO, O. A. E.; JONGEJAN, F. Simultaneous detection of *Anaplasma* and *Ehrlichia* species in ruminants and detection of *Ehrlichia ruminantium* in *Amblyomma variegatum* ticks by reverse line blot hybridization. **Veterinary Microbiology**, v. 89, p. 223-238, 2002.

- BELKHIA, H.; BEN SAID, M.; MABROUK, N.E.; SAIDANI, M.; CHERNI, C.; HASSEN, M.B.; BOUATTOUR, A.; MESSADI, L. Seasonal dynamics, spatial distribution and genetic analysis of *Anaplasma* species infecting small ruminants from Northern Tunisia. **Infection, Genetics and Evolution**, v. 54, p. 66-73, 2017.
- BEN SAID, M.; BELKHIA, H.; ALBERTI, A.; et al. Molecular Survey of *Anaplasma* Species in Small Ruminants. **Vector-Borne and Zoonotic Diseases**, v. 15, n. 10, p. 1–11, 2015.
- BEN SAID, M.; BELKHIA, H.; KARAOUD, M.; et al. First molecular survey of *Anaplasma bovis* in small ruminants from Tunisia. **Veterinary Microbiology**, v. 179, p. 322–326, 2015b.
- BEN SAID, M.; BELKHIA, H.; MABROUK, N. E.; et al. Molecular typing and diagnosis of *Anaplasma* spp. closely related to *Anaplasma phagocytophilum* in ruminants from Tunisia. **Ticks and Tick-borne Diseases**, v. 8, n. 3, p. 412–422, 2017.
- BIGUEZOTON, A.; NOEL, V.; ADEHAN, S.; et al. *Ehrlichia ruminantium* infects *Rhipicephalus microplus* in West Africa. **Parasites & Vectors**, n. 9, p. 4–7, 2016.
- BILGIC, H. B.; BAK, S.; KOSE, O.; et al. Prevalence of tick-borne haemoparasites in small ruminants in Turkey and diagnostic sensitivity of single-PCR and RLB. **Parasites & Vectors**, v. 10, p. 1–13, 2017.
- BIRKENHEUER, A.J.; LEVY, M.G.; BREITSCHWERDT, E.B. Development and evaluation of a seminested PCR for detection and differentiation of *Babesia gibsoni* (Asian genotype) and *B. canis* DNA in canine blood samples. **Journal of Clinical Microbiology**, v. 41, p. 4172-4177, 2003.
- BOES, K. M.; GONCAROV, K. O.; THOMPSON, C. A.; et al. Identification of a *Mycoplasma ovis*-like organism in a herd of farmed white-tailed deer (*Odocoileus virginianus*) in rural Indiana. **Veterinary Clinical Pathology**, v. 41, n. 1, p. 77–83, 2012.
- BONATO, L.; FIGUEIREDO, M.A.P.; GONÇALVES, L.R.; MACHADO, R.Z.; ANDRÉ, M.R. Occurrence and molecular characterization of *Bartonella* sp. and hemoplasmas in neotropical primates from Brazilian Amazon. **Comparative Immunology, Microbiology and Infectious Diseases**, n. 42, p. 15-20, 2015.
- BOSNOC, D.; BAREŠIĆ, M.; BRANIMIR, A.; SENTIĆ, M.; CEROVEC, M.; et al. Rare zoonosis (hemotropic mycoplasma infection) in a newly diagnosed systemic lupus erythematosus patient followed by a *Nocardia asteroides* pneumonia. **Brazilian Journal of Infectious Diseases**, p. 14, n. 1, p. 92-95, 2010.

- BRITO, D.R.B.; SANTOS, A.C.G.; GUERRA, R.M.S.N.C. Ectoparasitos em rebanhos de caprinos e ovinos na micro-região do Alto Mearim e Grajau, Estado do Maranhão. **Revista Brasileira de Parasitologia Veterinária**, v. 14, p. 59– 63, 2005.
- CARLSSON, H.E.; SCHAPIRO, S.J.; FARAH, I; HAU, J. Use of Primates in Research: A Global Overview. **American Journal of Primatology**, v. 63, p. 225-237, 2004.
- CEOLA, C.F.; SAMPAIO, J.; BLATT, S.L.; CORDOVA, C.M.M. *Mycoplasma* infection and inflammatory effects on laboratory rats bred for experimental research. **Revista Pan-Amazônica de Saúde**, v. 7, n. 2, p. 59-66, 2016.
- CHEN, Z.; LIU, Q.; FENG-CHAO, J.; et al. Detection of piroplasms infection in sheep, dogs and hedgehogs in Central China. **Infectious Diseases of Poverty**, v. 3, p. 18-25, 2012.
- CONRADO, F.O.; NASCIMENTO, N.C.; SANTOS, A.P.; ZIMPEL, C.K.; MESSICK, J.B.; et al. Occurrence and identification of hemotropic mycoplasmas (Hemoplasmas) in free ranging and laboratory rats (*Rattus norvegicus*) from two Brazilian zoos. **BMC Veterinary Research**, v. 11, p. 286-296, 2015.
- CONTAMIN, H.; MICHEL, J.C. Haemobartonellosis in Squirrel Monkeys (*Saimiri sciureus*): Antagonism between *Haemobartonella* sp. and Experimental *Plasmodium falciparum* Malaria. **Experimental Parasitology**, v. 91, p. 197-305, 1999.
- CUBILLA, M.P.; SANTOS, L.C.; MORAES, W.; CUBAS, Z.S.; LEUTENEGGER, M. et al. Occurrence of hemotropic mycoplasmas in non-human primates (*Alouatta caraya*, *Sapajus nigritus* and *Callithrix jacchus*) of southern Brazil. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 52, p. 6 – 13, 2017.
- DA SILVA, N.B.; TAUS, N.S.; JOHNSON, W.C.; MIRA, A.; et al. First report of *Anaplasma marginale* infection in goats, Brazil. **PLoS One**, v. 13, n. 8, 2018.
- DADDOW, K.N. The transmission of a sheep strain of *Eperythrozoon ovis* to goats and the development of a carrier state in the goats. **Australian Veterinary Journal**, v. 55, p. 605, 1979.
- DAHMANI, M.; MARIÉ, J.; SCANDOLA, P.; BRAH, S. *Anaplasma ovis* infects sheep in Niger. **Small Ruminant Research**, v. 151, p. 32–35, 2017.
- DARRIBA, D.; TABOADA, G.L.; DOALLO, R.; POSADA, D. jModelTest 2: more models, new heuristics and parallel computing. **Nature Methods**, v. 9, p. 772, 2012.
- DE LA FUENTE, J.; ATKINSON, M.W.; NARANJO, V.; MERA, I.G.F.; MANGOLD, A.J. Sequence analysis of the msp4 gene of *Anaplasma ovis* strains. **Veterinary Microbiology**, v. 119, n. 2-4, p. 375-381.

- DE LA FUENTE, J.; ATKINSON, M. W.; HOGG, J. T.; et al. Genetic Characterization of *Anaplasma ovis* Strains from Bighorn Sheep in Montana. **Journal of Wildlife Diseases**, v. 42, n. 2, p. 381–385, 2006.
- DE LA FUENTE, J.; TORINA, A.; CARACAPPA, S.; et al. Serologic and molecular characterization of *Anaplasma* species infection in farm animals and ticks from Sicily. **Veterinary Parasitology**, v. 133, p. 357–362, 2005.
- DERDAKOVA, M.; STEFANCIKOVA, A.; SPITALSKA, E.; et al. Emergence and genetic variability of *Anaplasma* species in small ruminants and ticks from Central Europe. **Veterinary Microbiology**, n. 153, p. 293-298, 2011.
- DESHUILLERS, P.L.; SANTOS, A.P.; NASCIMENTO, N.C.; HAMPEL, J.A.; BERGIN, I.L.; et al. Complete genome sequence of *Mycoplasma ovis* strain Michigan, a hemoplasma of sheep with two distinct 16S rRNA genes. **Genome Announcements**, v. 2, n. 1, p. 1-2, 2014.
- DILLBERGER, J.E.; LOUDY, D.E.; ADLER, R.R.; GASS, J.H. *Hemobartonella*-like Parasites in Cynomolgus Monkeys (*Macaca fascicularis*). **Veterinary Pathology**, v. 31, n. 3, p. 301-307, 1994.
- DUGAT, T.; CHASTAGNER, A.; LAGRÉE, A.; et al. A new multiple-locus variable-number tandem repeat analysis reveals different clusters for *Anaplasma phagocytophilum* circulating in domestic and wild ruminants. **Parasites & Vectors**, v. 7, p. 1–11, 2017.
- DUMLER, J. S.; BARBET, A. F.; BEKKER, C. P. J.; et al. Reorganization of genera in the families Rickettsiaceae and Anaplasmataceae in the order Rickettsiales: unification of some species of *Ehrlichia* with *Anaplasma*, *Cowdria* with *Ehrlichia* and *Ehrlichia* with *Neorickettsia*, descriptions of six new species combinations and designated of *Ehrlichia equi* ang ‘HGE agent’ as subjective synonyms of *Ehrlichia phagocytophila*. **International Journal of Systematic Evolutionary Microbiology**, v. 51, n. 6, p. 2145-2165, 2001.
- DYBVIG, K.; VOELKER, L.L. Molecular Biology of mycoplasmas. **Annual Review Microbiology**, v. 50, p. 25-27, 1996.
- EL IMAM, A.H.; HASSAN, S.M.; GAMEEL, A.A.; et al. Molecular identification of different *Theileria* and *Babesia* species infecting sheep in Sudan. **Annals of Parasitology**, v. 62, n. 2, p. 47-54, 2016.
- EMATER. Instituto Paranaense de Assistência Técnica e Extensão Rural. Projetos Ovinos e Caprinos, 2018. Disponível online em: <http://www.emater.pr.gov.br/modules/conteudo/conteudo.php?conteudo=72>. (Acessado em 04 de Novembro de 2018).

- EMBRAPA. Análise da PPM 2016: evolução dos rebanhos ovinos e caprios entre 2007 e 2016. Boletim do Centro de Inteligência e Mercado de Caprinos e Ovinos, n. 1, p. 1-13, 2017. Disponível online em: <https://ainfo.cnptia.embrapa.br/digital/bitstream/item/165640/1/CNPC-2017-Boletim-CIM.n1.pdf>. (Acessado em 04 de Novembro de 2018).
- ERSTER, O.; ROTH, A.; WOLLKOMIRSKY, R.; et al. Quantitative analysis of *Babesia ovis* infection in sheep and ticks. **Veterinary Parasitology**, v. 221, p. 39-45, 2016.
- FAEP. Federação da Agricultura do Estado do Paraná. Incentivo aos Produtores de Caprinos e Ovinos, 2017. Disponível online em: <http://sistemafaep.org.br/incentivo-aos-produtores-de-ovinos-e-caprinos/>. (Acessado em 04 de Novembro de 2018).
- FAO. Food and Agriculture Organization of the United Nations. FAOSTAT. Sheep. Live animals. Disponível online em: <http://www.fao.org/faostat/en/#search/sheep>. (Acessado em 10 de Dezembro de 2018).
- FERREIRA, E. P.; VIDOTTO, O.; ALMEIDA, J. C.; et al. Serological and molecular detection of *Theileria equi* in sport horses of northeastern Brazil. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 47, p. 72–76, 2016.
- GEBREKIDAN, H.; HAILU, A.; KASSAHUN, A.; ROHOU, I. *Theileria* infection in domestic ruminants in northern Ethiopia. **Veterinary Parasitology**, v. 200, p. 31–38, 2014.
- GHAH, R.R.; MUTINDA, M.; EZENWA, V.O. Limited sharing of tick-borne hemoparasites between sympatric wild and domestic ungulates. **Veterinary Parasitology**, n. 226, p. 167-173, 2016.
- GIANGASPERO, A.; MARANGI, M.; PAPINI, R.; et al. *Theileria* sp. OT3 and other tick-borne pathogens in sheep and ticks in Italy: Molecular characterization and phylogeny. **Ticks and Tick-borne Diseases**, v. 6, n. 1, p. 75–83, 2015.
- GIARDINI, N.D.; CHOCHLAKIS, D.; KRITSEPI-KONSTANTINOOU, M.; et al. Haemolytic disease in sheep attributed to a *Babesia lengau*-like organism. **Veterinary Record**, v. 170, n. 6, p. 155-157, 2012.
- GLASGOW, L.A.; MURRER, A.T.; LOMBARDI, P.S. *Eperythrozoon coccoides*. II. Effect on Interferon Production and Role of Humoral Antibody in Host Resistance. **Infection and Immunity**, p. 9, v. 2, p. 266-272, 1974.
- GLASGOW, L.A.; ODUGBEMI, T.; DWYER, P.; RITTERSON, A.L. *Eperythrozoon coccoides*. I. Effect on the Interferon Response in Mice. **Infection and Immunity**, v. 4, n. 4, p. 425-430.

- GOFF, W.; STILLER, D.; JESSUP, D.; et al. Characterization of an *Anaplasma ovis* Isolate from Desert Bighorn Sheep in Southern California Desert Sheep Bighorn. **Journal of Wildlife Diseases**, v. 29, n. 4, p. 540–546, 1993.
- GORDON, W. S.; BROWNLEE, A.; WILSON, D. R.; MACLEOD, J. Tick-Borne Fever: A Hitherto Undescribed Disease of Sheep. **Journal of Comparative Pathology and Therapeutics**, v. 45, n. 2, p. 301–307, 1932.
- GRAZZIOTIN, A. L.; DUARTE, J. M. B.; SZABÓ, M. P. J.; et al. Prevalence and Molecular Characterization of *Mycoplasma ovis* in Selected Free-Ranging Brazilian Deer Populations. **Journal of Wildlife Diseases**, v. 47, n. 4, p. 1005–1011, 2011.
- GRAZZIOTIN, A. L.; SANTOS, A. P.; GUIMARAES, A. M. S.; et al. *Mycoplasma ovis* in captive cervids: Prevalence, molecular characterization and phylogeny. **Veterinary Microbiology**, v. 152, n. 3–4, p. 415–419, 2011b.
- GUIMARÃES, J.H.; BATTESTI, D.M.B.; TUCCI, E.C. **Ectoparasitos de importância veterinária**. 1a Edição, São Paulo: Editora Plêiade, 2001.
- HAGHI, S.M.M.; FAKHAR, M.; SHARIF, M.; et al. Molecular Identification of Ovine *Babesia* sp. in North Iran. **Research in Molecular Medicine**, v. 1, n. 1, p. 35-39, 2013.
- HAMPEL, J. A.; SPATH, S. N.; BERGIN, I. L.; et al. Prevalence and Diagnosis of Hemotrophic *Mycoplasma* Infection in Research Sheep and Its Effects on Hematology Variables and Erythrocyte Membrane Fragility. **Comparative Medicine**, v. 64, n. 6, p. 478–485, 2014.
- HASHEMINISAB, S.S.; MORADI, P.; WRIGHT, I. A four-year epidemiological and chemotherapy survey of babesiosis and theileriosis, and tick vectors in sheep, cattle and goats in Dehghan, Iran. **Annals of Parasitology**, v. 64, n. 1, p. 43-48, 2018.
- HASSAN, M.A.; RAOOFI, A.; LOTFOLLAHZADEH, S.; JAVANBANKHT, J. Clinical and cytological characteristics and prognostic implications on sheep and goat *Theileria* infection in north of Iran. **Journal of Parasitic Diseases** v. 39, n. 2, p. 190-193, 2015.
- HEIDAPOUR-BAMI, M.; HADDADZADEH, H.R.; KAZEMI, B.; et al. Molecular identification of ovine *Theileria* species by a new PCR-RFLP method. **Veterinary Parasitology**, v. 12, n. 3-4, p. 171-177, 2009.
- HOELZLE, K.; WINKLER, M.; KRAMER, M.M.; WITTENBRINK, M.M.; DIECKMANN, S.M.; HOELZLE, L.E. Detection of *Candidatus Mycoplasma haemobos* in cattle with anemia. **Veterinary Journal**, v. 187, p. 408-410, 2011.

- HOOSHMAND-RAD, P.; HAWA, N. J. Malignant Theileriosis of Sheep. **Tropical Animal Health and Production**, v. 5, p. 97–102, 1973.
- HORNOK, S.; HAJTÓS, I.; MELI, M.L.; FARKAS, I.; GÖNCZI, E. First molecular identification of *Mycoplasma ovis* and *Candidatus M. haemoovis* from goat, with lack of haemoplasma PCR-positivity in lice. **Acta Veterinaria Hungarica**, v. 60, p. 355–360, 2012.
- HORNOK, N.; HOFF, B.; GANTER, M. *Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an epierythrocytic agent of haemolytic anemia in sheep and goats. **International Journal of Systematic and Evolutionary Microbiology**, v. 54, p. 365–71, 2004.
- HORNOK, S.; ELEK, V.; DE LA FUENTE, J.; NARANJO, V.; FARKAS, R.; MAJOROS, G.; FOLDVÁRI, G. First serological and molecular evidence of endemicity of *Anaplasma ovis* and *A. marginale* in Hungary. **Veterinary Microbiology**, v. 122, p. 316–322, 2007.
- HORNOK, S.; MELI, M. L.; ERDOS, A.; et al. Molecular characterization of two different strains of haemotropic mycoplasmas from a sheep flock with fatal haemolytic anaemia and concomitant *Anaplasma ovis* infection. **Veterinary Microbiology**, v. 136, p. 372–377, 2009.
- HORNOK, S.; SUGAR, L.; FERNÁNDEZ, I.F.; et al. Tick- and fly-borne bacteria in ungulates: the prevalence of *Anaplasma phagocytophilum*, haemoplasmas and rickettsiae in water buffalo and deer species in Central Europe, Hungary. **BMC Veterinary Research**, v. 14, p. 98–104, 2018.
- HORTA, S.; BARRETO, M.C.; PEPE, A.; CAMPOS, J.; OLIVA, A. Highly sensitivity method for diagnosis of subclinical *B. ovis* infection. **Ticks and Tick-Borne Diseases**, v. 5, p. 902–906, 2014.
- HSU, D.Y.M.; GEIMAN, Q.M. Synergistic effect of *Haemobartonella muris* on *Plasmodium berghei* in white rats. **American Journal of Tropical Medicine and Hygiene**, v. 1, n. 5, p. 747–760, 1952.
- IBGE. Pesquisa da Pecuária Municipal. Efetivos de rebanho, por tipo de rebanho, 2017. Disponível online em: <https://sidra.ibge.gov.br/tabela/3939>. (Acessado em 22 de Julho de 2018).
- Institute of Medicine and National Research Council. Use of Laboratory Animals in Biomedical and Behavioral Research. Washington, DC, Estados Unidos: The National Academies Press, 1988.
- JALALI, R. Molecular detection and identification of *Anaplasma* species in sheep from Ahvaz, Iran. **Iranian Journal of Veterinary Research**, v. 14, n. 1, p. 50–56, 2013.

- JIA, N.; ZHENG, Y.; JIANG, J.; et al. Human babesiosis caused by a novel *Babesia crassa*-like pathogen: A case series. **Clinical Infectious Diseases**, v. 67, n. 7, p. 110–1119, 2018.
- JIAXUN, L.; HONG, Y. Theileriosis of Sheep and Goats in China. **Tropical Animal Health Production**, v. 29, p. 85-105, 1997.
- JOHNSON, K. A.; NASCIMENTO, N.C.; BAUER, A. E.; WENG, H.; et al. Detection of hemoplasma infection of goats by use of a quantitative polymerase chain reaction assay and risk factor analysis for infection. **American Journal of Veterinary Research**, v. 77, n. 8, 2016.
- KATOH, K.; STANDLEY, D.M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. **Molecular Biology Evolution**, v. 30, n. 4, p. 772–780, 2013.
- KOCAN, K. M. et al. The natural history of *Anaplasma marginale*. **Veterinary Parasitology**, v. 167, n. 2-4, p. 95-107, 2010.
- KOCAN, K.M.; STILLER, D. Development of *Anaplasma ovis* (Rickettsiales: Anaplasmataceae) in male *Dermacentor andersoni* (Acari: Ixodidae) transferred from infected to susceptible sheep. **Journal of Medical Entomology**, v. 29, n. 1, p. 98-107, 1991.
- KUTTLER, A. K. L. *Anaplasma* Infections in Wild and Domestic Ruminants: A Review. **Journal of Wildlife Diseases**, v. 20, n. 1, p. 12–20, 1984.
- LADBURY, G. A. F.; STUEN, S.; THOMAS, R.; et al. Dynamic Transmission of Numerous *Anaplasma phagocytophilum* Genotypes among Lambs in an Infected Sheep Flock in an Area of Anaplasmosis Endemicity. **Journal of Clinical Microbiology**, v. 46, n. 5, p. 1686–1691, 2008.
- LALOY, E.; PETIT, E.; BOULOUIS, H.; et al. Dynamics of natural infection by *Anaplasma phagocytophilum* in a dairy cattle herd in Brittany, France. **Clinical Microbiology and Infection**, v. 15, p. 24–25, 2009.
- LANE, J.; JUBB, T.; SHEPHARD, R.; WEBB-WARE, J.; FORDYCE, G. MLA Final Report: Priority List of Endemic Diseases for the Red Meat Industries. Meat and Livestock Australia. Sydney, Australia (2015). Disponível em: <https://www.mla.com.au/research-and-development/search-rd-reports/final-report-details/Animal-Health-and-Biosecurity/Priority-list-of-endemic-diseases-for-the-red-meat-industries/2895>. (Acessado em 13 de Março de 2019).
- LI, H.; ZHENG, Y.; MA, L.; et al. Human infection with a novel tick-borne *Anaplasma* species in China: a surveillance study. **The Lancet Infectious Diseases**, v. 15, n. 6, p. 663–670, 2015.

- LI, Y.; GUAN, G.; MA, M.; LIU, J.; REN, Q.; LUO, J.; YIN, H. *Theileria ovis* discovered in China. **Experimental Parasitology**, n. 127, p. 304-307, 2011.
- LIU, A. H.; YIN, H.; GUAN, G. Q.; et al. At least two genetically distinct large *Babesia* species infective to sheep and goats in China. **Veterinary Parasitology**, v. 147, p. 246-251, 2007.
- MA, M.; CHEN, Z.; LIU, A.; REN, Q.; LIU, J.; et al. Biological parameters of *Rhipicephalus* (*Boophilus*) *microplus* (Acari: Ixodidae) fed on rabbits, sheep, and cattle. **Korean Journal of Parasitology**, v. 54, p. 301-305, 2016.
- MACHADO, C. A. L.; VIDOTTO, O.; CONRADO, F. O.; et al. *Mycoplasma ovis* infection in goat farms from northeastern Brazil. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 55, n. July, p. 1-5, 2017.
- MACHADO, R. Z.; MARIA, M.; TEIXEIRA, G.; et al. Molecular diagnosis and genetic diversity of tick-borne Anaplasmatidae agents infecting the African buffalo *Syncerus caffer* from Marromeu Reserve in Mozambique. **Parasites & Vectors**, v. 9, p. 454-463, 2016.
- MAGGI, R.G.; MASCARELLI, P.E.; BALAKRISHNAN, N.; ROHDE, C.M.; KELLY, C.M.; et al. “*Candidatus Mycoplasma haemomacae*” and *Bartonella quintana* bacteremia in cynomolgus monkeys. **Journal of Clinical Microbiology**, v. 51, n. 5, p. 1408-1411, 2013.
- MARIANTE, A.S.; MCMANUS, C.; MENDONÇA, J.F. **Country Report on the State of Animal Genetics Resources Brazil**. 1^o Edição. Brasília: Embrapa Genetic Resources and Biotechnology, 2003.
- MARTINEZ-HERNÁNDEZ, J. M.; BALLADOS-GONZÁLEZ, G. G.; FERNÁNDEZ-BANDALA, D.; et al. Molecular detection of *Mycoplasma ovis* in an outbreak of hemolytic anemia in sheep from Veracruz, Mexico. **Tropical Animal Health and Production**, v. 51, n. 1, p. 243-248/
- MASON, R.W.; STATHAM, P. The determination of the level of *Eperythrozoon ovis* parasitaemia in chronically infected sheep and its significance to the spread of infection. **Animal Health Laboratory**, v. 68, n. 3, p. 115-116, 1991.
- MASON, R.W.; STATHAM, P. Susceptibility of sheep and goats to *Eperythrozoon ovis* infection. **Australian Veterinary Journal**, v. 68, p. 116-117, 1991.
- MCMANUS, C.; PAIVA, S. R.; ARAÚJO, R. O. DE. Genetics and breeding of sheep in Brazil. *Genética e melhoramento de ovinos no Brasil*. **Revista Brasileira de Zootecnia**, v.39, p. 236-246, 2010.

- MELO-JUNIOR, O.A.; ALBERNAZ, A.P.; THIEBAUT, J.T.L. et al. Ocorrência de *Anaplasma bovis* (Donatien & Lestoquard, 1936, Dumler et al., 2001) na região de Bom Jesus do Itabapoana, RJ. **Ciência Animal Brasileira**, v. 11, n. 4, p.888-8891, 2010.
- MELTEZER, M.I.; PERRY, B.D.; DONACHIE, P.L. Mortality percentages related to heartwater and the economic impact of heartwater disease on large-scale commercial farms in Zimbabwe. **Preventive Veterinary Medicine**, v. 26, p. 187-199, 1996.
- MESSICK, J. B. Hemotrophic mycoplasmas (hemoplasmas): a review and new insights into pathogenic potential. **Veterinary Clinical Pathology**, v. 33, n. 1, p. 20–22, 2004.
- MGHIRBI, Y.; ROS-GARCIA, A.; IRIBAR, P.; et al. A molecular study of tick-borne haemoprotozoan parasites (*Theileria* and *Babesia*) in small ruminants in Northern Tunisia. **Veterinary Parasitology**, v. 198, p. 72-77, 2013.
- MOLYNEUX, D.H. Relationship between *Eperythrozoon coccoides* and *Trypanosoma* (Trypanozoon) *brucei brucei* in experimentally infected mice. **Annals of Tropical Medicine & Parasitology**, v. 64, n. 3, p. 325-328, 1970.
- MONGRUEL, A. C. B.; BENEVENUTE, J. L.; IKEDA, P.; et al. Detection of *Anaplasma* sp. phylogenetically related to *A. phagocytophilum* in a free-living bird in Brazil. **Revista Brasileira de Parasitologia Veterinária**, v. 26, n. 4, p. 505–510, 2017.
- MORZARIA, S.P. Theileriosis. In: DELVES, P.J; ROITT, I. **Encyclopedia of Immunology**. São Diego, CA, Estados Unidos: Academic Press, 1998.
- NAGORE, D.; GARCÍA-SANMARTÍN, J.; GARCÍA-PÉREZ, A. L.; et al. Identification, genetic diversity and prevalence of *Theileria* and *Babesia* species in a sheep population from Northern Spain q. **International Journal for Parasitology**, v. 34, p. 1059–1067, 2004.
- NEIMARK, H.; HOFF, B.; GANTER, M. *Mycoplasma ovis* comb. nov. (formerly *Eperythrozoon ovis*), an epierythrocytic agent of haemolytic anaemia in sheep and goats. **International Journal of Systematic and Evolutionary Microbiology**, v. 54, p. 365–371, 2004.
- NEIMARK, H.; JOHANSSON, K.; RIKIHISA, Y.; et al. Proposal to transfer some members of the genera *Haemobartonella* and *Eperythrozoon* to the genus *Mycoplasma* with descriptions of ‘*Candidatus Mycoplasma haemofelis*’, ‘*Candidatus Mycoplasma haemomuris*’, ‘*Candidatus Mycoplasma haemosuis*’ and ‘*Candidatus*. **International Journal of Systematic and Evolutionary Microbiology**, v. 51, p. 891–899, 2001.

NEITZ, W. O.; ALEXANDER, R. A.; DU TOIT, P. J. *Eperythrozoon ovis* (sp. nov.) Infection in Sheep. **Onderstepoort Journal of Veterinary Science and Animal Industry**, v. 3, n. 2, p. 263–271, 1934.

NEITZ, W.O. Tick-borne diseases as a hazard in the rearing of calves in Africa. **Bulletin de l'Office International des Épidémiologies**, v. 62, p. 607-625, 1964.

NELSON, J.B. An ascites tumor appearing during the passage of *Eperythrozoon coccoides* in mice. **Journal of Experimental Medicine**, v. 103, n. 6, p. 743-752, 1956.

OCHIRKHUU, N.; KONNAI, S.; ODBILEG, R.; et al. Molecular Epidemiology Survey and Genetic Characterization of Anaplasma Species in Mongolian Livestock. **Vector Borne and Zoonotic Diseases**, n. 17, v. 8, p. 539-549, 2017.

OGDEN, N. H.; CASEY, A. N. J.; FRENCH, N. P.; et al. Natural *Ehrlichia phagocytophila* transmission coefficients from sheep 'carriers' to *Ixodes ricinus* ticks vary with the numbers of feeding ticks. **Parasitology**, v. 124, p. 127–136, 2002.

OHTAKE, Y.; NISHIZAWA, I.; SATO, M.; WATANABE, Y.; NISHIMURA, T. *Mycoplasma ovis* Detected in Free-Living Japanese Serows, *Capricornis crispus*. **Journal of Veterinary Medicine and Science**, v. 73, n. 3, p. 371–373, 2011.

OTT, K.J.; ASTIN, J.K.; LESLIE, A.S. *Eperythrozoon coccoides* and Rodent Malaria: *Plasmodium chabaudi* and *Plasmodium berghei*. **Experimental Parasitology**, v. 21, p. 68-77, 1967.

OZUBEK, S.; AKTAS, M. Molecular and Parasitological Survey of Ovine Piroplasmosis, Including the First Report of *Theileria annulata* (Apicomplexa: Theileridae) in Sheep and Goats from Turkey. **Journal of Medical Entomology Advance**, v. 54, n. 1, p.212-220, 2016.

PAIVA, D. F.; SILVÉRIO, V.C.; PAIVA, D.A.F.; MCMANUS, C.; et al. Origin of the Main Locally Adapted Sheep Breeds of Brazil: A RFLP-PCR Molecular Analysis. **Archivos de Zootecnia**, v. 54, p. 395-399, 2005.

Peters W, Molyneux DH, Howells RE: *Eperythrozoon* and *Haemobartonella* in monkeys. **Ann Trop Med Parasitol** 1974; 68: 47-50.

MAGGI, R.G.; COMPTON, S.M.; TRULL, C.L.; MASCARELLI, P.E.; MOZAYENI, B.R.; et al. Infection with hemotropic *Mycoplasma* species in patients with or without extensive arthropod or animal contact. **Journal of Clinical Microbiology**, v. 51, p. 3237–3241, 2013.

RAJENHRAN, J.; GUNASEKARAN, P. Microbial phylogeny and diversity: Small subunit ribosomal RNA sequence analysis and beyond. **Microbiology Research**, v. 166, p. 99-110, 2011.

- RAMALHO, A.C., GUERRA, R.R.; MONGRUEL, A.C.B.; VIDOTTO, O.; LUCENA, R.B.; et al. *Mycoplasma* sp. infection in captive Marcgrave's capuchin monkeys (*Sapajus flavius*). **Comparative Immunology, Microbiology and Infectious Diseases**, v. 51, p. 34-36, 2017.
- RAZIN, S.; YOGEV, D.; NAOT, Y. Molecular Biology and Pathogenicity of Mycoplasmas. **Microbiology Molecular Biology Review**, v. 62, n. 4, p. 1094-1156, 1998.
- RENNEKER, S.; ABDO, J.; BAKHEIT, M.; et al. Coinfection of Sheep with *Anaplasma*, *Theileria* and *Babesia* Species in the Kurdistan Region, Iraq. **Transboundary and Emerging Diseases**, v. 60, b. 2, p. 113-118, 2013b.
- RENNEKER, S.; ABDO, J.; SALIH, D. E. A.; KARAGENC, T. Can *Anaplasma ovis* in Small Ruminants be Neglected any Longer? **Transboundary and Emerging Diseases**, v. 60, p. 105–112, 2013.
- RIKIHISA, Y.; KAWAHARA, M.; WEN, B.; KOCIBA, G.; FUERST, P.; et al. Western Immunoblot Analysis of *Haemobartonella muris* and Comparison of the 16S rRNA Gene Sequences of *H. muris*, *H. felis* and *Eperythrozoon suis*. **Journal of Clinical Microbiology**, v. 35, n. 4, p. 823-829, 1997.
- RIKIHISA, Y. Mechanisms of Obligatory Intracellular Infection with *Anaplasma phagocytophilum*. **Clinical Microbiology Reviews**, v. 24, n. 3, p. 469–489, 2011.
- RIKIHISA, Y.; KAWAHARA, M.; WEN, B.; et al. Western Immunoblot Analysis of *Haemobartonella muris* and Comparison of 16S rRNA Gene Sequences of *H. muris*, *H. felis*, and *Eperythrozoon suis*. **Journal of Clinical Microbiology**, v. 35, n. 4, p. 823–829, 1997.
- RINGO, A. E.; MOUMOUNI, P. F. A.; TAIÖE, M.; et al. Molecular analysis of tick-borne protozoan and rickettsial pathogens in small ruminants from two South African provinces. **Parasitology International**, v. 67, n. 2, p. 144–149, 2018.
- RJEIBI, M. R.; DARGHOUTH, M. A.; REKIK, M.; et al. First Molecular Identification and Genetic Characterization of *Theileria lestoquardi* in Sheep of the Maghreb Region. **Transboundary and Emerging Diseases**, v. 63, p. 278–284, 2016b.
- RJEIBI, M. R.; DARGHOUTH, M.; GHARBI, M. Prevalence of *Theileria* and *Babesia* species in Tunisian sheep. **Onderstepoort Journal of Veterinary Research**, v. 83, n. 1, p. a1040, 2016.
- ROBINSON, W.F.; ROBINSON, N.A. Chapter 1 – Cardiovascular System. In: JUBB, K.V.F.; KENNEDY, P.C.; PALMER, N. **Pathology of domestic animals**. São Diego, CA, Estados Unidos: Academic Press, 1993. p. 1-101.

- ROS-GARCIA, A.; BARANDIKA, J.F.; GARCÍA-PEREZ, A.L.; JUSTE, R.A.; HURTADO, A. Assessment of exposure to piroplasms in sheep grazing in communal mountain pastures by using a multiplex DNA bead-based suspension array. **Parasites & Vectors**, v. 6, p. 277-288, 2013.
- SANTOS, A.P.; SANTOS, R.P.; BIONDO, A.W.; DORA, J.M.; GOLDANI, L.Z.; et al. Hemoplasma infection in HIV-positive Patient, Brazil. **Emerging Infectious Diseases**, v. 14, n. 12, p. 1922-1924, 2008.
- SASHIDA, H.; SUZUKI, Y.; ROKUHARA, S.; NAGAI, K.; HARASAWA R. Molecular Demonstration of Hemotropic *Mycoplasmas* in Wild Japanese Monkeys (*Macaca fuscata*). **Journal of Veterinary Medicine Science**, v. 76, n. 1, p. 97-101, 2014.
- SCHNITTGER, L.; YIN, H.; GUBBELS, M. J.; et al. Phylogeny of sheep and goat *Theileria* and *Babesia* parasites. **Parasitology Research**, v. 91, p. 398–406, 2003.
- SELA, I.; ASHKENAZY, H.; KATOH, K.; PUPKO, T. GUIDANCE2: accurate detection of unreliable alignment regions accounting for the uncertainty of multiple parameters. **Nucleic Acid Research**, v. 43, p. W7-W14, 2015.
- SEVINC, F.; ZHOU, M.; CAO, S.; et al. Haemoparasitic agents associated with ovine babesiosis: A possible negative interaction between *Babesia ovis* and *Theileria ovis*. **Veterinary Parasitology**, v. 252, n. January, p. 143–147, 2018.
- SHABANA, I. I.; ALHADLAG, N. M.; ZARAKET, H. Diagnostic tools of caprine and ovine anaplasmosis: a direct comparative study. **BMC Veterinary Research**, v.14, p. 1–8, 2018.
- SHAHZAD, W.; NOOR, H.; AHMAD, M.; et al. Prevalence and Molecular Diagnosis of *Babesia ovis* and *Theileria ovis* in Lori Sheep at Livestock Experiment Station (LES), Bahadurnagar, Okara, Pakistan. **Iranian Journal of Parasitology**, v. 8, n. 4, p. 570-578, 2013.
- Sheriff D, Clapp KH, Reid MA. *Eperythrozoon ovis* infection in south Australia. *Aust Vet J* 1966; 42: 169.
- SHERIFF, D.; CLAPP, K. H.; REID, M. *Eperythrozoon ovis* INFECTION IN SOUTH AUSTRALIA. **Australian Veterinary Journal**, v. 42, p. 169, 1966.
- SILVEIRA, A. W.; OLIVEIRA, G. G.; SANTOS, L. M.; et al. Natural infection of the South America Tapir (*Tapirus terrestris*) by *Theileria equi*. **Journal of Wildlife Diseases**, v. 53, n. 2, p. 411–413, 2017.
- SILVEIRA, J. A. G.; OLIVEIRA, C. H. S. DE; SILVESTRE, B. T.; et al. Molecular assays reveal the presence of *Theileria* spp. and *Babesia* spp. in Asian water buffaloes (*Bubalus*

bubalis, Linnaeus, 1758) in the Amazon region of Brazil. **Ticks and Tick-borne Diseases**, v. 7, n. 5, p. 1017–1023, 2016.

STUEN, S. Tick-Borne Fever (*Anaplasma phagocytophilum* Infection) in Sheep – A Review. **Journal of Veterinary Medicine and Research**, v. 3, 2016.

STUEN, S.; OKSTAD, W.; SAGEN, A. M. Intrauterine Transmission of *Anaplasma phagocytophilum* in Persistently Infected Lambs. **Veterinary Sciences**, v. 5, p.25-32, 2018.

SYKES, J.E.; LINDSAY, L.L.; MAGGI, R.G.; BREITSCHWERDT, E. Human Coinfection with *Bartonella henselae* and Two Hemotropic *Mycoplasma* Variants Resembling *Mycoplasma ovis*. **Journal of Clinical Microbiology**, v. 48, n. 10, p. 3782-3785, 2010.

TABAEI, S.J.S.; SPOTIN, A.; POURIRAN, R.; et al. Molecular characterization of *Theileria* spp. in livestock and the first report on the occurrence of *Theileria* sp. OT3 in Iran. **Acta Parasitologica**, v. 63, n. 3, p. 515–521, 2018.

TAGAWA, M.; TAKEUCHI, T.; FUJISAWA, T.; et al. A Clinical Case of Severe Anemia in a Sheep Coinfected with *Mycoplasma ovis* and ‘*Candidatus* Mycoplasma haemovis’ in Hokkaido, Japan. **Journal of Veterinary Medicine and Science**, v. 74, n. 1, p. 99-102, 2012.

TAGELDIN, M. H.; FADIYA, A. A.; SABRA, A. A.; ISMAILY, S. I. A. Theileriosis in sheep and goats in the Sultanate of Oman. **Tropical Animal Health and Production**, v. 37, p. 491–493, 2005.

TAYLOR, M.A.; COOP, R.L.; WALL, R.L. **Veterinary Parasitology**. 3a Edição, Iowa, Estados Unidos: Wiley-Blackwell, 2007.

THOMAS, R. J.; BIRTLES, R. J.; RADFORD, A. D.; WOLDEHIWET, Z. Recurrent Bacteraemia in Sheep Infected Persistently with *Anaplasma phagocytophilum*. **Journal of Comparative Pathology**, v. 147, n. 2–3, p. 360–367, 2012.

TIAN, Z.; LIU, G.; YIN, H.; et al. Parasitology International First report on the occurrence of *Theileria* sp. OT3 in China. **Parasitology International**, v. 63, n. 2, p. 403–407, 2014.

TORINA, A.; GALINDO, R. C.; VICENTE, J.; et al. Characterization of *Anaplasma phagocytophilum* and *A. ovis* infection in a naturally infected sheep flock with poor health condition. **Tropical Animal Health and Production**, v. 42, p. 1327–1331, 2010.

TRATCHENBERG, S. Quick Guide: Mollicutes. **Current Biology**, v. 15, n. 12, p. PR483-484, 2005.

UENO, H.; GONÇALVES, P.C. **Manual para Diagnóstico das Helmintoses de Ruminantes**. 4a Edição, Tóquio: Japan International Cooperation Agency, 1998.

- UILENBERG, G. *Babesia* - A historical overview. **Veterinary Parasitology**, v. 138, p. 3–10, 2006.
- VIEIRA, T. S. W. L.; QUROLLO, B. A.; MONGRUEL, A. C. B.; et al. Potentially Same Novel *Ehrlichia* Species in Horses in Nicaragua and Brazil. **Emerging Infectious Diseases**, v. 24, n. 5, p. 180238, 2018.
- VIERIA, M. I. B.; COSTA, M. M.; OLIVEIRA, M. T.; et al. Serological detection and molecular characterization of piroplasmids in equids in Brazil. **Acta Tropica**, v. 179, p. 81-87, 2018b.
- VILHENA, H.; TVARIJONAVICIUTE, A.; CÉRON, J.J.; RAMIRO PASTORINHO, M.; MARTINEZ-SUBIELA, S.; et al. Acute phase proteins response in cats naturally infected by hemotropic mycoplasmas. **Comparative Immunology, Microbiology and Infectious Diseases**, v. 56, p. 1-5, 2018.
- WANG, X.; CUI, Y.; ZHANG, Y.; SHI, K.; YAN, Y.; et al. Molecular characterization of hemotropic mycoplasmas (*Mycoplasma ovis* and ‘*Candidatus Mycoplasma haemovis*’) in sheep and goats in China. **BMC Veterinary Research**, v. 13, p. 142-150, 2017.
- WANG, J.; ZHANG, Y.; CUI, Y. et al. A rapid, simple and sensitive loop-mediated isothermal amplification method to detect *Anaplasma bovis* in sheep and goat samples. **Parasitology International**, n. 67, p. 70-73, 2018.
- WEISS, D.J.; WARDROP, K.J.; SCHALM, O.W. **Schalm’s Veterinary Hematology**. 4a Edição, Ames, Iowa, Estados Unidos: Wiley-Blackwell, 2010.
- WOLDEHIWET, Z. The natural history of *Anaplasma phagocytophilum*. **Veterinary Parasitology**, v. 167, p. 108-122, 2010.
- WOODS, J.E.; BREWER, M.M.; HAWLEY, J.R.; MISNEWSKI, N.; LAPPIN, M.R. Evaluation of experimental transmission of *Candidatus Mycoplasma haemominutum* and *Mycoplasma haemofelis* by *Ctenocephalides felis* to cats. **American Journal of Veterinary Research**, v. 66, n. 6, p. 1008-1012, 2005.
- YANG, J.; HAN, R.; NIU, Q.; et al. Ticks and Tick-borne Diseases Occurrence of four *Anaplasma* species with veterinary and public health significance in sheep, northwestern China. **Ticks and Tick-borne Diseases**, v. 9, n. 1, p. 82–85, 2018.
- YANG, J.; LI, Y.; LIU, Z.; et al. Molecular detection and characterization of *Anaplasma* spp. in sheep and cattle from Xinjiang, northwest China. **Parasites & Vectors**, v. 8, p. 108–115, 2015.

- YASINI, S.; KHAKI, Z.; RAHBARI, S.; KAZEMI, B. Hematologic and Clinical Aspects of Experimental Ovine Anaplasmosis Caused by *Anaplasma ovis* in Iran. **Iranian Journal of Parasitology**, v. 7, n. 4, p. 91-98, 2012.
- YERUHAM, I.; HADANI, A.; GALKER, F. Some epizootiological and clinical aspects of ovine babesiosis caused by *Babesia ovis* — a review. **Veterinary Parasitology**, v. 74, p. 153-163, 1998.
- YIN, H.; SCHNITTGER, L.; LUO, J.; SEITZER, U.; AHMED, J. S. Ovine theileriosis in China: a new look at an old story. **Parasitology Research**, v. 101, n. 2, p. 191–195, 2007.
- YOUSEFI, A.; RAHBARI, S.; SHAYAN, P. et al. Molecular detection of *Anaplasma marginale* and *Anaplasma ovis* in sheep and goat in west highland pasture of Iran. **Asian Pacific Journal of Tropical Biomedicine**, v. 7, n. 5, p. 455- 459, 2017.
- ZAEEMI, M.; HADDADZADEH, H. Identification of different *Theileria* species (*Theileria lestoquardi*, *Theileria ovis*, and *Theileria annulata*) in naturally infected sheep using nested PCR – RFLP. **Parasitology Research**, v. 108, p. 837–843, 2011.
- ZHAN, L.; CAO, W.; JIANG, J.; et al. *Anaplasma phagocytophilum* from Rodents and Sheep, China. **Emerging Infectious Diseases**, v. 16, n. 5, 2010.
- ZHANG, J.; KELLY, P.; LI, J.; et al. Molecular detection of *Theileria* spp. in Livestock in Caribbean Islands. **BioMed Research International**, v. 8, p. 1-8, 2015.
- ZHANG, Y.; LV, Y.; CUI, Y.; et al. First molecular evidence for the presence of *Anaplasma* DNA in milk from sheep and goats in China. **Parasitology Research**, v. 115, n. 7, p. 1–7, 2016b.
- ZHANG, Y.; LV, Y.; ZHANG, F.; ZHANG, W. Molecular and phylogenetic analysis of *Anaplasma* spp. in sheep and goats from six provinces of China. **Journal of Veterinary Science**, v. 17, n. 4, p. 523-529, 2016.
- ZOBBA, R.; ANFOSSI, A. G.; VISCO, S.; et al. Ticks and Tick-borne Diseases Cell tropism and molecular epidemiology of *Anaplasma platy* -like strains in cats. **Ticks and Tick-borne Diseases**, v. 6, n. 3, p. 272–280, 2015.
- ZYGOYIANNIS, D. Sheep production in the world and in Greece. **Small Ruminant Research**, v. 62, p. 143–147, 2006.